Amphora by sherd count per period. Figure is the % of that form found in that period

FORMS	0	II	II-III	III	III-IV	IV	V	VI	VII
ABUNG	14.29%	0.00%	0.00%	28.57%	0.00%	42.86%	7.14%	7.14%	0.00%
Beltran 1	16.67%	45.83%	4.17%	17.50%	0.83%	5.00%	7.50%	1.67%	0.83%
Dr 20	16.32%	4.40%	2.54%	48.08%	1.80%	13.52%	9.99%	3.35%	0.00%
Dr 2-4	11.40%	71.05%	3.51%	5.70%	0.44%	4.82%	1.75%	0.44%	0.88%
Dr1 or 2-4	18.98%	59.12%	5.11%	7.30%	2.19%	5.84%	0.73%	0.73%	0.00%
Dressel 1	4.06%	87.65%	2.03%	4.23%	0.00%	1.10%	0.51%	0.42%	0.00%
Gauloise	31.28%	9.48%	0.00%	26.54%	5.21%	14.22%	10.90%	2.37%	0.00%
Haltern 70	33.33%	16.67%	0.00%	50.00%	0.00%	0.00%	0.00%	0.00%	0.00%
Kapitan II	0.00%	0.00%	0.00%	0.00%	0.00%	100.00%	0.00%	0.00%	0.00%
Pascual 1	13.16%	52.63%	7.89%	10.53%	0.00%	5.26%	5.26%	5.26%	0.00%
AMISC	20.32%	22.46%	0.00%	20.32%	2.67%	18.72%	10.70%	4.28%	0.53%

Amphora by sherd count by period: Figure is the % of the total assemblage of that period represented by that form.

FORMS	0	II	II-III	III	III-IV	IV	V	VI	VII
ABUNG	0.40%	0.00%	0.00%	0.41%	0.00%	1.81%	0.44%	1.27%	0.00%
Beltran 1	4.03%	3.70%	5.68%	2.16%	2.00%	1.81%	3.96%	2.53%	25.00%
Dr 20	53.02%	4.77%	46.59%	79.57%	58.00%	65.86%	70.93%	68.35%	0.00%
Dr 2-4	5.24%	10.89%	9.09%	1.33%	2.00%	3.32%	1.76%	1.27%	50.00%
Dr1 or 2-4	5.24%	5.44%	7.95%	1.03%	6.00%	2.42%	0.44%	1.27%	0.00%
Dressel 1	9.68%	69.62%	27.27%	5.13%	0.00%	3.93%	2.64%	6.33%	0.00%
Gauloise	13.31%	1.34%	0.00%	5.75%	22.00%	9.06%	10.13%	6.33%	0.00%
Haltern 70	0.40%	0.07%	0.00%	0.31%	0.00%	0.00%	0.00%	0.00%	0.00%
Kapitan II	0.00%	0.00%	0.00%	0.00%	0.00%	0.60%	0.00%	0.00%	0.00%
Pascual 1	1.01%	1.34%	3.41%	0.41%	0.00%	0.60%	0.88%	2.53%	0.00%
AMISC	7.66%	2.82%	0.00%	3.90%	10.00%	10.57%	8.81%	10.13%	25.00%

Amphora by sherd weight per period. Figure is the % of that form found in that period VII II Ш III-IV IV VI **FORMS** 0 II-III V **ABUNG** 12.90% 0.00% 0.00% 54.84% 0.00%16.13% 0.00% 16.13% 0.00% Beltran 1 11.20% 50.71% 8.64% 19.45% 0.18%4.32% 0.00% 4.77% 0.73% Dr 20 14.91% 3.44% 2.07% 45.69% 1.39% 14.74% 8.64% 9.13% 0.00% Dr 2-4 11.71% 66.07% 5.35% 6.95% 3.01% 1.29% 4.59% 0.33% 0.71% 8.17%Dr1 or 2-4 21.60% 51.69% 9.03% 2.74% 4.04% 0.35% 2.37% 0.00% Dr1A 71.18% 28.82% 0.00%0.00%0.00%0.00%0.00%0.00%0.00% Dressel 1 1.47% 0.08% 0.58% 0.00% 16.56% 74.62% 3.97% 0.00% 2.73% Gauloise 23.75% 6.19% 0.00% 41.52% 2.21% 9.56% 5.92% 10.86% 0.00% 0.00% Haltern 70 42.24% 11.97% 0.00% 45.79% 0.00% 0.00% 0.00% 0.00% Kapitan II 0.00%0.00%0.00%0.00%0.00%100.00 0.00%0.00%0.00% % 5.57% Pascual 1 9.23% 54.64% 13.34% 11.69% 0.00% 0.00% 5.52% 0.00% 21.97% 20.97% 0.00% 22.85% 9.80% 8.19% 0.26% **AMISC** 1.50% 14.46% all 16.10% 34.68% 2.43% 26.38% 0.89% 9.32% 4.70% 5.45% 0.05%

Amphora by sherd weight by period: Figure is the % of the total assemblage of that period represented by that form.

FORMS 0 II II-III III III-IV IV V VI

FORMS	0	II	II-III	III	III-IV	IV	V	VI	VII
ABUNG	0.01%	0.00%	0.00%	0.02%	0.00%	0.02%	0.00%	0.04%	0.00%
Beltran 1	1.92%	4.05%	9.84%	2.04%	0.56%	1.28%	0.00%	2.42%	40.31%
Dr 20	44.21%	4.73%	40.72%	82.68%	74.48%	75.52%	87.72%	79.96%	0.00%
Dr 2-4	2.33%	6.09%	6.03%	0.65%	1.17%	2.38%	2.04%	0.76%	45.74%
Dr1 or 2-4	5.19%	5.76%	14.37%	1.20%	11.89%	1.68%	0.29%	1.68%	0.00%
Dr1A	0.98%	0.18%	0.00%	0.00%	0.00%	0.00%	0.00%	0.00%	0.00%
Dressel 1	35.66%	74.59%	20.94%	5.22%	0.00%	10.16%	0.56%	3.67%	0.00%
Gauloise	4.39%	0.53%	0.00%	4.68%	7.35%	3.05%	3.74%	5.92%	0.00%
Haltern 70	0.77%	0.10%	0.00%	0.51%	0.00%	0.00%	0.00%	0.00%	0.00%
Kapitan II	0.00%	0.00%	0.00%	0.00%	0.00%	0.82%	0.00%	0.00%	0.00%
Pascual 1	0.85%	2.32%	8.10%	0.65%	0.00%	0.88%	0.00%	1.49%	0.00%
AMISC	3.70%	1.64%	0.00%	2.35%	4.54%	4.20%	5.65%	4.07%	13.95%
all	100.00%	100.00%	100.00%	100.00%	100.00%	100.00%	100.00%	100.00%	100.00%