

Palynological studies at Sand, Applecross

Fraser M. Green and Kevin J. Edwards¹

(Department of Geography and Environment, University of Aberdeen)

Introduction

Material collected in two small kubiena tins from Trench A by Dr Cressey was examined for its organic and palynological content. The first of these, 'Sand 1' was of particular interest as it contained the transition from an inferred midden deposit to a minerogenic soil. 'Sand 3' was taken from lower in the profile where minerogenic soil graded into a friable sandstone unit. Both samples originated from the east-facing sections of the trench, excavated beneath a loose sandstone slab of obscure provenance.

Description

Sand 1 consisted of a 10 cm thickness of poorly consolidated material. The upper 3 cm contained the inferred midden material and had numerous inclusions of shell in a sandy matrix. The lower 7 cm, in terms of visibility to the eye and within the sieve retent seemed to be composed entirely of very sandy soil. Sand 3 consisted of very sandy soil throughout.

Methods and presentation of results

Both sections were sampled contiguously in 1 cm thicknesses and prepared for pollen analysis using standard NaOH, HF and acetolysis techniques (Faegri and Iversen, 1989) after sieving through a 180 μ m mesh. Carbonate

¹ Department of Geography and Environment, University of Aberdeen, Elphinstone Road, Aberdeen, AB24 3UF
f.m.green@abdn.ac.uk
kevin.edwards@abdn.ac.uk

tablets containing spores of *Lycopodium clavatum* were added at the start of sample preparation, to enable estimates of palynomorph and charcoal concentrations (Stockmarr, 1971). Samples were then mounted on slides using silicone oil. Replicate samples for both sections were analysed for loss on ignition (LOI) for 16 hours at 360 °C, and were mixed with deionised water for pH tests. The latter were performed because of indications of effervescence from HCl in the shelly samples (top 3 cm of Sand 1).

The pollen and spore content for both samples was negligible, while microscopic charcoal was more frequent. A large number of microscope slides was searched for each level. As a result of this sparseness, the palynomorphs were individually noted and identified instead (pollen type nomenclature follows Stace [1997], amended after Bennett [1994]). Microscopic charcoal was sufficiently abundant for counts of fragments greater than 400 μm^2 size, and involved counting individual fragments in two dimensions using a microscope eye-piece micrometer. The values for charcoal are expressed as concentrations ($\text{mm}^2 \text{cm}^{-3}$). With the exception of some shell fragments in the upper 3 cm of Sand 1, the bulk of the sieve retent was minerogenic material. The data displayed in Figures 1 and 2 was processed using TILIA and TILIA.GRAPH (Grimm, 1991).

Discussion

The results for Sand 1 (Figure 1) illustrate the highly minerogenic nature of the material, the organic content remaining consistently low and generally less than 4 % LOI throughout the profile. Even though effervescence in HCl was evident only during the pretreatment of the shelly samples, the pH of all samples was remarkably high throughout, perhaps as a consequence of leaching of dissolved shell material rich in CaCO_3 , or from lime-rich superficial deposits found locally (country rock is acid Torridonian Sandstone). These very alkaline conditions would explain the lack of palynological evidence, given that pollen is typically only preserved when the pH is less than 5.5 (Dimpleby 1985). In spite of this, it was decided to persist with pollen and

spore analysis because the method has produced interpretable results in some calcareous contexts (Dimbleby and Evans 1974). Microscopic charcoal, though of significant concentration, varies greatly throughout the section, although the supposed midden segment has enhanced charcoal values. If this is a result of the inclusion of hearth waste, the lack of macroscopic charcoal is surprising and would argue against the presence of an immediately adjacent domestic fire. Microscope analysis of the larger shell fragments from the top 3 cm revealed morphological features suggesting a mix of cockle and limpet.

The palynological data were extremely limited and only *Corylus avellana*-type (cf. hazel), Poaceae (grasses), Cyperaceae (sedges), *Myriophyllum verticillatum* (whorled water-milfoil), Pteropsida (monolete) indet. (undifferentiated ferns) and *Sphagnum* (bog moss) were present. All these plants save one are part of the flora today. *Myriophyllum verticillatum*, an aquatic plant of basic fresh water, seems to be absent from Scotland now (Perring and Walters 1990) and was perhaps so in the past (cf. Godwin 1975).

Unidentified spore A (Figure 1) was relatively abundant. This spore is spherical in all views, well preserved, inaperturate, and 35-45 μm in diameter. The exine is very thick (5 μm) and is dominated by strong sculptural elements – a network of high relief ridges forming a chaotic web. The lumina between the ridges occupy 40-60 % of the grain surface and are of irregular shape. The grain appears to be tectate (in that it possesses a foot layer) but lacks columellae. Its absence from the pollen and spore keys suggests a possible pre-Quaternary origin with derivation from local drift materials. It is interesting to note however, that spore A is not present in the upper levels of Sand 3, despite this deposit reportedly representing the bottom part of the minerogenic material described in Sand 1.

Figure 2 shows the results for Sand 3. The organic content in the upper part of the profile matches that of the lower section of Sand 1, but quickly reduces as the transition to an even sandier matrix takes place. This is reflected in the charcoal curve which decreases significantly in the basal 4 cm. As seen in

Sand 1 the pH is consistently high, despite the lack of shell material in these samples, with a corresponding lack of palynomorphs. This would again suggest that the parent material for the soils was highly alkaline.

Sand 3 is of even less palynological interest than Sand 1, producing only one identifiable grain of *Corylus avellana*-type pollen. Unidentified spore B is very different from the example in Sand 1. It too is spherical and inaperturate, but is half the size (18-25 μm) and relatively featureless. The exine is thin and almost universally psilate, though some palynomorphs show a faint ridging pattern on their surface, producing sunken areas of a polygonal pattern, visible only under $\times 1000$ magnification. An estimated 10-16 of these polygons cover the entire grain surface, if present at all. It has been suggested to us that these relatively featureless palynomorphs may be the spores of a liverwort or obscure moss (Dr Andrew McMullen, pers. comm.).

Acknowledgement

Mrs Maureen Lamb and Dr Andrew McMullen are thanked for providing laboratory assistance and guidance.

References

- Bennett, K.D. (1994). *Annotated catalogue of pollen and pteridophyte spore types of the British Isles*. <http://www.kv.geo.uu.se/pc-intro.html>.
- Dimbleby, G.W. (1985). *The palynology of archaeological sites*. London: Academic Press.
- Dimbleby, G.W. and Evans, J.G. (1974). Pollen and land-snail analysis of calcareous soils. *Journal of Archaeological Science* 1, 117-133.
- Faegri, K. and Iversen, J. (1989). *Textbook of pollen analysis*. 4th edition by K. Faegri, P.E. Kaland and K. Krzywinski. Chichester: John Wiley & Sons.
- Godwin, H. (1975). *History of the British flora: a factual basis for phytogeography*. 2nd edition. Cambridge: Cambridge University Press.
- Grimm, E.C. (1991). *TILIA and TILIA.GRAPH*. Springfield: Illinois State Museum.
- Perring, F.H. and Walters, S.M. (1990). *Atlas of the British flora*. Botanical Society of the British Isles.
- Stace, C. (1997). *New flora of the British Isles*. 2nd edition. Cambridge: Cambridge University Press.
- Stockmarr, J. (1971). Tablets with spores used in absolute pollen analysis. *Pollen et Spores* 13, 615-621.




Figure 1. Palynological, loss on ignition and pH data from Sand 1. Charcoal concentrations are in values of mm² cm⁻³ and palynomorph numbers represent raw counts.




Figure 2. Palynological, loss on ignition and pH data from Sand 3. Charcoal concentrations are in values of $\text{mm}^2 \text{cm}^{-3}$ and palynomorph numbers represent raw counts.