ISSN 1749-8775

## KING JOHN'S HUNTING LODGE, LACOCK, WILTSHIRE TREE-RING ANALYSIS OF TIMBERS

SCIENTIFIC DATING REPORT

Matt Hurford, Martin Bridge and Cathy Tyers





ARCHAEOLOGICAL SCIENCE Research Department Report Series 101-2010

## KING JOHN'S HUNTING LODGE, LACOCK, WILTSHIRE

## TREE-RING ANALYSIS OF TIMBERS

Matt Hurford, Dr Martin Bridge and Cathy Tyers

NGR: ST 9166 6857

© English Heritage

ISSN 1749-8775

The Research Department Report Series incorporates reports from all the specialist teams within the English Heritage Research Department: Archaeological Science; Archaeological Archives; Historic Interiors Research and Conservation; Archaeological Projects; Aerial Survey and Investigation; Archaeological Survey and Investigation; Archaeological Survey of London. It replaces the former Centre for Archaeology Reports Series, the Archaeological Investigation Report Series and the Architectural Investigation Report Series.

Many of these are interim reports which make available the results of specialist investigations in advance of full publication. They are not usually subject to external refereeing, and their conclusions may sometimes have to be modified in the light of information not available at the time of the investigation. Where no final project report is available, readers must consult the author before citing these reports in any publication. Opinions expressed in Research Department reports are those of the author(s) and are not necessarily those of English Heritage.

Requests for further hard copies, after the initial print run, can be made by emailing: Res.reports@english-heritage.org.uk or by writing to:

Or Dy Writing LO: English Haritago Fort Cur

English Heritage, Fort Cumberland, Fort Cumberland Road, Eastney, Portsmouth PO4 9LD Please note that a charge will be made to cover printing and postage.

#### SUMMARY

Dendrochronological analysis was undertaken on 11 samples taken from the extant remains of the medieval open-hall core of King John's Hunting Lodge. This resulted in the production of a single site chronology, LCKJSQ01 which comprises nine samples with an overall length of 171 rings dating to the years AD 1148–1318. The results identified that the dated timbers used in the construction of the open-hall probably represent a single programme of felling in the AD 1320s.

#### CONTRIBUTORS

Matt Hurford, Dr Martin Bridge, and Cathy Tyers

#### ACKNOWLEDGEMENTS

The laboratories would like to thank Mrs Margaret Vaughan, the manager of King John's Hunting Lodge, and Graham Heard, National Trust property manager for Lacock, for giving permission to undertake the work. Thanks are also due to Avis Lloyd for arranging access and for providing additional background material on the building and to Clive Carter of the Wiltshire Buildings Record for providing drawings. The Wiltshire Buildings Record project work was supported by the English Heritage Historic Environment Enabling Programme. The dendrochronological work was funded by English Heritage and co-ordinated by Peter Marshall from the English Heritage Scientific Dating Team.

#### **ARCHIVE LOCATION**

Wiltshire Archaeological Service The Wiltshire and Swindon History Centre Cocklebury Road Chippenham SN15 3QN

#### DATE OF INVESTIGATION

2009-10

#### CONTACT DETAILS

Matt Hurford and Cathy Tyers Dendrochronology Laboratory Graduate School of Archaeology University of Sheffield West Court, 2 Mappin Street Sheffield SI 4DT Tel: 0114 276 3146 Email: m.hurford@sheffield.ac.uk, c.m.tyers@sheffield.ac.uk

Dr Martin Bridge UCL Institute of Archaeology 31–34 Gordon Square London WC1H 0PY Email: martin.bridge@ucl.ac.uk

## CONTENTS

Introduction I
King John's Hunting LodgeI
Sampling2
Analysis and Results
Interpretation4
Discussion and Conclusion4
Bibliography6
Tables7
Figures
Data of Measured Samples
Appendix: Tree-Ring Dating20
The Principles of Tree-Ring Dating
The Practice of Tree-Ring Dating at the Nottingham Tree-Ring Dating Laboratory
I. Inspecting the Building and Sampling the Timbers
2. Measuring Ring Widths25
3. Cross-matching and Dating the Samples25
4. Estimating the Felling Date
5. Estimating the Date of Construction
6. Master Chronological Sequences
7. Ring Width Indices
References

## INTRODUCTION

In 2009 the Wiltshire Buildings Record successfully obtained support through the English Heritage Historic Environment Enabling Programme for their project 'Wiltshire cruck buildings and other archaic roof types'. The detailed aims and objectives of the project are set out in the Project Design (Lloyd 2009). The overall aim is to establish a typological chronology of archaic roof types and hence elucidate the development of carpentry techniques in the county. This will then facilitate detailed comparison with other counties allowing Wiltshire to be placed in the regional context. Investigation of these late-medieval buildings (c AD 1200 – c AD 1550) will combine building survey, historical research, and dendrochronological analysis.

A series of buildings identified by the Wiltshire Buildings Record as having the potential to contribute to the aims and objectives of the project was assessed for dendrochronological suitability during 2009. In order to maximise the potential, these detailed dendrochronological assessments and the WBR's assessments of the significance of the buildings within the project informed the selection of the buildings subsequently subjected to detailed study.

A single final report produced by the Wiltshire Buildings Record (forthcoming a) will summarise the overall results from the project. However, each building included in the project will have an associated individual report produced by the WBR (forthcoming b), whilst the primary archive of the dendrochronological analysis is the English Heritage Research Department Report Series.

A brief introduction to dendrochronology can be found in the Appendix. However, further details can be found in the guidelines published by English Heritage (1998) which are also available on the English Heritage website (http://www.english-heritage.org.uk/publications/dendrochronology-guidelines/).

## King John's Hunting Lodge

This grade II\* listed building is located immediately to the west of St. Cyriac's Church in Lacock (ST 9166 6857; Figs 1 and 2) with its north elevation fronting onto Church Street. The following information is summarised from the Wiltshire Buildings Record report (forthcoming b) on King John's Hunting Lodge and the listed building record (http://lbonline.english-heritage.org.uk).

The earliest extant fabric, thought to be thirteenth-century in date, survives from an earlier building. This includes a single two-centred arched doorway, originally one of a pair, and a series of timbers that were retained as part of the screens passage in the presumed early fourteenth-century rebuilding of the house with a two-bay open hall (Fig 3). The house was extended to the rear with the addition of the south-east wing (Fig 4), probably during the early eighteenth century. The encasing of the building in rubble stone may also date to this period.

The focus of this investigation was primarily on the extant remains of the open hall structure comprising two trusses, A and B (Figs 5 and 6). The presence of an incomplete plain-chamfered windbrace extending westwards of truss A, towards a postmedieval softwood truss, implies that this bay was also part of the medieval open hall, which is therefore thought to have originally comprised three bays. This screens passage and open-hall structure were of raised-cruck construction. Trusses A and B both have slightly cranked collars; truss A was originally a double-collar truss, though the lower collar is no longer extant. There is a single pair of plain-chamfered windbraces in bay 2 and the extant arch braces have hollow-chamfer mouldings with run-out stops. There is a single row of purlins and a diagonally set ridge piece. Both trusses have saddles but on truss A an apex block above the saddle takes the ridge piece whereas on truss B the saddle is notched to take the ridge piece directly. Ten original common rafters are present, five in each of the two extant bays. The opposing pairs of common rafters between trusses A and B have original side-lapped and pegged fixings for a former louvre. Smoke-blackening of the timbers associated with the original open-hall house is apparent throughout the roof space.

The dendrochronological potential of the south-east wing was also investigated at this stage, as this had not been feasible during the initial pre-sampling assessment of the building due to access being severely restricted. The roof is of two bays and incorporates three trusses. It is a principal-rafter roof with tiebeams and threaded purlins (Fig 7). The principals are connected with a staggered haunch joint at the apex and there is a diagonally set ridge piece. Numerous original common rafters survive. The tiebeam was visible on northernmost of these trusses (Fig 7) and in the gable wall truss; both of these trusses also had a number of original studs.

#### SAMPLING

The assessment of the timbers in the south-east wing indicated that these were all elm (*Ulmus* spp.) and hence would have been excluded from this project regardless of date. Sampling and analysis by tree-ring dating of the timbers associated with the medieval open hall were commissioned by English Heritage. It was hoped to provide independent dating evidence for the construction of this element of the building and hence inform the overall objectives of the *Wiltshire cruck buildings and other archaic roof types* project. The dendrochronological study also formed part of the English Heritage-funded training programme for the first author.

A total of 13 oak (*Quercus* spp.) timbers associated with the extant remains of the medieval open hall were sampled by coring. Each sample was given the code LCK-J (for Lacock, King John's) and numbered 01–13. The sampling encompassed as wide a range of elements as possible, whilst focussing on those timbers with the best dendrochronological potential. Unfortunately no samples were taken from the common rafters in Bay 1, as they were derived from fast-grown trees and were considered highly unlikely to provide samples with sufficient numbers of rings for reliable dendrochronological analysis.

The location of the samples was noted at the time of coring and marked on the drawings subsequently provided by the Wiltshire Buildings Record, these being reproduced here as Figures 8–10. Further details relating to the samples can be found in Table 1. In this table the timbers have been located and numbered following the scheme on the drawings provided with the trusses and bays being labelled from west to east.

## ANALYSIS AND RESULTS

Each of the 13 samples was prepared by sanding and polishing. It was seen at this point that one sample, LCK-J03, had too few rings for reliable dating and one, LCK-J05, was elm, and so these were rejected from this programme of analysis. The annual growth rings of the remaining 11 samples were measured, the data of these measurements being given at the end of this report.

The ring sequences derived from these 11 samples were initially compared with each other by the Litton/Zainodin grouping procedure (see Appendix) allowing a single group of nine series to be formed, the samples of this group cross-matching with each other as shown in the bar diagram (Fig 11). The analytical process was aided by the use of software written by Tyers (2004).

The nine grouped series were combined at their indicated offsets to form site chronology LCKJSQ01. Intra-site cross-matching (see below) indicated the possibility that some timbers may have been derived from the same tree as suggested by *t*-values in excess of 10.0. However, to maintain consistency between all of the dendrochronological reports on individual buildings within this project, these potential same-tree series were not combined prior to incorporation into the site chronology, thus following the Nottingham Tree-Ring Dating Laboratory standard practice. This site master was then compared to an extensive range of reference chronologies for oak, indicating repeated cross-matching when the first ring in the sequence is AD 1148 and the last ring date is AD 1318. The evidence for this is given in Table 2.

The site chronology was compared with the remaining two ungrouped samples but there was no further satisfactory cross-matching. The two ungrouped samples were then compared individually with the reference chronologies, but there was no conclusive cross-matching and these samples must, therefore, remain undated.

The analysis can be summarised as follows:

Site chronology LCKJSQ01	Number of samples 9	Number of rings 171	Date span (where dated) AD 1148–1318
	2		undated
	2		unmeasured

## INTERPRETATION

For consistency the sapwood estimate used in all of the dendrochronological reports on individual buildings within this project is the Nottingham Tree-Ring Dating Laboratory estimate of 15-40 (95% confidence) rings. This is used to calculate felling date ranges for samples with incomplete sapwood or felled-after dates for samples which are heartwood only.

The medieval core of King John's Hunting Lodge is represented by nine dated timbers associated with the roof structure (Fig 11). A full complement of sapwood was present on one of the dated timbers, LCK-J11, but due to its highly friable state, approximately 1–5mm of sapwood, including the bark edge, was lost during sample preparation prior to measurement. The average ring width of this entire sequence is 0.69mm (Table 1) but the overall growth trend is towards narrower rings, the outer 10 measured rings having an average ring width of 0.51mm. This suggests that approximately 2–10 rings have been lost. Thus, with a last measured ring date of AD 1318, it is likely that the felling date is in the AD 1320s.

Two of the other dated samples retained their heartwood/sapwood boundary ring. Sample LCK-J01 has a heartwood/sapwood boundary ring of AD 1307 and hence an estimated felling date in the range of AD 1322–1347. Sample LCK-J02 has a heartwood/sapwood boundary ring of AD 1290 and hence an estimated felling date in the range of AD 1305–1330. However, allowing for the outermost measured ring, this can be truncated to AD 1319–30.

The remaining six dated samples in site chronology LCKJSQ01 have no trace of sapwood and thus it is not possible to calculate their likely felling date ranges. Five of these samples have final-ring dates within six years of each other, ranging from AD 1301 to AD 1307, with the remaining one having a last-ring date of AD 1276.

All the dated samples are therefore clearly broadly coeval. The three retaining sapwood or heartwood/sapwood boundary have overlapping felling dates or felling date ranges and hence could represent a single felling, although the variation in heartwood/sapwood date could suggest that they represent a single programme of felling that may have spanned a number of years in the AD 1320s. This interpretation is supported by the overall level of cross-matching between the individual series (Table 3) which includes some *t*-values sufficiently high (>10) to suggest the possibility that the timbers could have been derived from a single tree.

## DISCUSSION AND CONCLUSION

Tree-ring analysis has confirmed that the extant remains of the former three-bay openhall core of King John's Hunting Lodge is of early fourteenth-century date, as previously thought based on other evidence. The trees used in the construction of the open hall roof appear likely to have been felled as part of a single programme of felling in the AD I 320s, with construction following shortly afterwards. The dated timbers represent both extant trusses and the intervening bay, bay 2. It is unfortunate that no timbers could be dated from bay 1, with the common rafters proving to have too few rings and the ridge proving to be elm. However, given the structural integrity of the remains and the consistent level of smoke blackening, it appears likely, but unproven by dendrochronological analysis, that the timbers present in bay 1 are coeval with those from the rest of the roof. However, it cannot be entirely ruled out that they could be reused timbers salvaged from the thirteenth-century structure.

The high overall level of intra-site cross-matching, with many of the *t*-values in excess of 5.0 (Table 3) suggests that these timbers probably originated from the same woodland source. The average ring widths of the timbers analysed varies from 0.65 to 1.71mm, suggesting that some of the timbers were derived from quite dense woodland containing slow-growing trees. The site chronology, LCKJSQ01, generally produces the highest *t*-values, and thus shows the greatest degree of similarity, with reference chronologies from Wiltshire and the surrounding counties (Table 2). This suggests that it is likely that the timbers were derived from a relatively local woodland source.

### BIBLIOGRAPHY

Bridge, M C, 2001 Tree-ring analysis of timbers from the Abbey Barn, Glastonbury, Somerset, Centre for Archaeol Rep, **39/2001** 

Bridge, M C, 2002 Tree-ring analysis of timbers from Meare Manor Farmhouse, St Mary's Road, Meare, Somerset, Centre for Archaeol Rep, 103/2002

Groves, C, 2005 Dendrochronological research in Devon: phase 1, Centre for Archaeol Rep, 56/2005

English Heritage, 1998 Dendrochronology: guidelines on producing and interpreting dendrochronological dates, London

Howard, R E, Laxton, R R, and Litton, C D, 2001 Tree-ring analysis of timbers from Exeter Cathedral, Exeter, Devon -part I, Centre for Archaeol Rep, **47/2001** 

Hurford, M, Bridge, M, and Tyers, C, forthcoming a Wick Farm Cottage, Heddington Wick, Wiltshire: tree-ring analysis of timbers, English Heritage Res Dep Rep Ser

Hurford, M, Bridge, M, and Tyers, C, forthcoming b Dauntsey House, Dauntsey, Wiltshire: tree-ring analysis of timbers,, English Heritage Res Dep Rep Ser

Lloyd, A, 2009 Wiltshire cruck buildings and other archaic roof types: an archaeological and dendrochronological analysis of medieval timber construction in the county, EH Historic Environment Enabling Programme No 5104 Project Design

Siebenlist-Kerner, V, Schove, D, and Fletcher, J M, 1978 The barn at Great Coxwell, Berkshire, in *Dendrochronology in Europe* (ed J M Fletcher), BAR Int Ser, **51**, 295–320

Tyers, I, 2003 Tree-ring analysis of oak timbers from the South Transept and Nave roofs of the Church of St John the Baptist, Bradworthy, Devon, Centre for Archaeol Rep, 2/2003

Tyers, I, 2004 Dendro for Windows program guide, 3rd edn, ARCUS Rep, 500b

Wiltshire Buildings Record forthcoming a Wiltshire cruck buildings and other archaic roof types: an archaeological analysis of medieval timber construction in the county, WBR report

Wiltshire Buildings Record forthcoming b King John's Hunting Lodge, Lacock, WBR report

## TABLES

Sample number	Sample location	Total rings	Sapwood rings	Average ring width (mm)	Cross-section dimensions (mm)	First measured ring date (AD)	Last heartwood ring date (AD)	Last measured ring date (AD)
LCK-J01	Truss A collar	152	h/s	1.00	160×320	1156	1307	1307
LCK-J02	Truss A north principal	170	28	1.27	150×280	1149	1290	1318
LCK-J03	Truss A south principal	nm			250×300			
LCK-J04	Truss A saddle	157	no h/s	1.71	150×340	1148		1304
LCK-J05	Bay I ridge - elm	nm			120×130			
LCK-J06	Bay 2 south common rafter I	128	no h/s	0.91	90×110	1180		1307
LCK-J07	Bay 2 north common rafter I	56	h/s	1.43	90×100			
LCK-J08	Bay 2 north common rafter 2	54	no h/s	0.75	90×90			
LCK-J09	Bay 2 north common rafter 3	78	no h/s	0.65	90×100	1224		1301
LCK-J10	Bay 2 north common rafter 4	86	no h/s	1.07	90×100	9		1276
LCK-JII	Truss B north principal	120	24c (1–5mm	0.69	140×220	1199	1294	1318
			sap lost)					
LCK-J12	Truss B south principal	147	no h/s	1.22	140×380	1160		1306
LCK-J13	Truss B collar	82	no h/s	1.06	150×310	1226		1307

Table 1: Details of tree-ring samples from King John's Hunting Lodge, Lacock, Wiltshire

nm = not measured

h/s = the heartwood/sapwood ring is the last ring on the sample

c = complete sapwood was present on the timber but some was lost during coring/sample preparation

Table 2: Results of the cross-matching of site sequence LCKJSQ01 and relevant reference chronologies when the first-ring date is AD 1148 and the last-	
ring date is AD 1318	

Reference chronology	t-value	Span of chronology	Reference
Abbey Barn, Glastonbury, Somerset	10.0	AD 1095-1334	(Bridge 2001)
Exeter Cathedral, Exeter, Devon	9.9	AD 1132-1315	(Howard et al 2001)
Wick Farm Cottage, Heddington Wick, Wiltshire	9.4	AD 1158-1335	(Hurford et al forthcoming a)
Dauntsey House, Dauntsey, Wiltshire	9.1	AD 1122-1355	(Hurford et al forthcoming b)
Rudge, Morchard Bishop, Devon	9.0	AD 1124-1315	(Groves 2005)
Manor Farmhouse, Meare, Somerset	8.8	AD 1156-1315	(Bridge 2002)
Church of St John the Baptist, Bradworthy, Devon	8.8	AD 1125-1367	(Tyers 2003)
Great Coxwell barn, Oxfordshire	8.8	AD 1043-1267	(Siebenlist-Kerner et al 1978)

## $\infty$ Table 3: Matrix showing the t-values obtained between the individual ring sequences in site chronology LCKJSQ01; - indicates t-values less than 3.00

	lck-j02	lck-j04	lck-j06	lck-j09	lck-j10	lck-j l l	lck-j12	lck-j I 3
lck-j01	5.16	8.20	9.20	7.79	3.22	6.36	5.46	10.65
lck-j02		6.57	4.28	-	3.91	5.23	5.74	3.53
lck-j04			8.13	4.27	3.64	3.97	5.48	7.04
lck-j06				7.68	4.02	5.75	3.10	7.95
lck-j09					3.74	4.24	3.30	8.60
lck-j10						-	3.17	3.01
lck-j l l							10.17	4.62
lck-j12								3.02

## FIGURES



Figure 1: Map to show the location of King John's Hunting Lodge, Lacock, Wiltshire (based on the Ordnance Survey map with permission of the Controller of Her Majesty's Stationery Office, © Crown Copyright)




Figure 2: Map to show the location of King John's Hunting Lodge, Lacock, Wiltshire (based on the Ordnance Survey map with permission of the Controller of Her Majesty's Stationery Office, © Crown Copyright




Figure 3: The north elevation of King John's Hunting Lodge viewed looking south-west showing the approximate truss locations



Figure 4: The south-east wing gable of King John's Hunting Lodge viewed looking north



Figure 5: East face of truss A



Figure 6: West face of truss B



Figure 7: North face of the northernmost truss of the south-east wing




Figure 8: East face of truss A




Figure 9: West face of truss B





Figure 10: Sample location plan (later replacement rafters are shown dashed in)



c= complete sapwood exists on the timber but part of the sapwood on the sample has been lost during preperation

Figure 11: Bar diagram of the samples in site chronology LCKJSQ01

#### DATA OF MEASURED SAMPLES

Measurements in 0.01mm units

## APPENDIX: TREE-RING DATING

#### The Principles of Tree-Ring Dating

Tree-ring dating, or dendrochronology as it is known, is discussed in some detail in the Laboratory's Monograph, An East Midlands Master Tree-Ring Chronology and its uses for dating Vernacular Building (Laxton and Litton 1988) and Dendrochronology: Guidelines on Producing and Interpreting Dendrochronological Dates (English Heritage 1988). Here we will give the bare outlines. Each year an oak tree grows an extra ring on the outside of its trunk and all its branches just inside its bark. The width of this annual ring depends largely on the weather during the growing season, about April to October, and possibly also on the weather during the previous year. Good growing seasons give rise to relatively wide rings, poor ones to very narrow rings and average ones to relatively average ring widths. Since the climate is so variable from year to year, almost randomlike, the widths of these rings will also appear random-like in sequence, reflecting the seasons. This is illustrated in Figure A1 where, for example, the widest rings appear at irregular intervals. This is the key to dating by tree rings, or rather, by their widths. Records of the average ring widths for oaks, one for each year for the last 1000 years or more, are available for different areas. These are called master chronologies. Because of the random-like nature of these sequences of widths, there is usually only one position at which a sequence of ring widths from a sample of oak timber with at least 70 rings will match a master. This will date the timber and, in particular, the last ring.

If the bark is still on the sample, as in Figure A1, then the date of the last ring will be the date of felling of the oak from which it was cut. There is much evidence that in medieval times oaks cut down for building purposes were used almost immediately, usually within the year or so (Rackham 1976). Hence if bark is present on several main timbers in a building, none of which appear reused or are later insertions, and if they all have the same date for their last ring, then we can be quite confident that this is the date of construction or soon after. If there is no bark on the sample, then we have to make an estimate of the felling date; how this is done is explained below.

# The Practice of Tree-Ring Dating at the Nottingham Tree-Ring Dating Laboratory

1. Inspecting the Building and Sampling the Timbers. Together with a building historian the timbers in a building are inspected to try to ensure that those sampled are not reused or later insertions. Sampling is almost always done by coring into the timber, which has the great advantage that we can sample *in situ* timbers and those judged best to give the date of construction, or phase of construction if there is more than one in the building. The timbers to be sampled are also inspected to see how many rings they have. We normally look for timbers with at least 70 rings, and preferably more. With fewer rings than this, 50 for example, sequences of widths become difficult to match to a unique position within a master sequence of ring widths


and so are difficult to date (Litton and Zainodin 1991). The cross-section of the rafter shown in Figure A2 has about 120 rings; about 20 of which are sapwood rings – the lighter rings on the outside. Similarly the core has just over 100 rings with a few sapwood rings.

To ensure that we are getting the date of the building as a whole, or the whole of a phase of construction if there is more than one, about 8–10 samples per phase are usually taken. Sometimes we take many more, especially if the construction is complicated. One reason for taking so many samples is that, in general, some will fail to give a date. There may be many reasons why a particular sequence of ring widths from a sample of timber fails to give a date even though others from the same building do. For example, a particular tree may have grown in an odd ecological niche, so odd indeed that the widths of its rings were determined by factors other than the local climate! In such circumstances it will be impossible to date a timber from this tree using the master sequence whose widths, we can assume, were predominantly determined by the local climate at the time.

Sampling is done by coring into the timber with a hollow corer attached to an electric drill and usually from its outer rings inwards towards where the centre of the tree, the pith, is judged to be. An illustration of a core is shown in Figure A2; it is about 150mm long and 10mm diameter. Great care has to be taken to ensure that as few as possible of the outer rings are lost in coring. This can be difficult as these outer rings are often very soft (see below on sapwood). Each sample is given a code which identifies uniquely which timber it comes from, which building it is from and where the building is located. For example, CRO-A06 is the sixth core taken from the first building (A) sampled by the Laboratory in Cropwell Bishop. Where it came from in that building will be shown in the sampling records and drawings. No structural damage is done to any timbers by coring, nor does it weaken them.

During the initial inspection of the building and its timbers the dendrochronologist may come to the conclusion that, as far as can be judged, none of the timbers have sufficient rings in them for dating purposes and may advise against sampling to save further unwarranted expense.

All sampling by the Laboratory is undertaken according to current Health and Safety Standards. The Laboratory's dendrochronologists are insured.



on the outside just inside the bark. The year of each ring can be determined by counting back from the outside ring, which grew in 1976



Figure A2: Cross-section of a rafter, showing sapwood rings in the left-hand corner, the arrow points to the heartwood/sapwood boundary (H/S); and a core with sapwood; again the arrow is pointing to the H/S. The core is about the size of a pencil



Figure A3: Measuring ring widths under a microscope. The microscope is fixed while the sample is on a moving platform. The total sequence of widths is measured twice to ensure that an error has not been made. This type of apparatus is needed to process a large number of samples on a regular basis



Figure A4: Three cores from timbers in a building. They come from trees growing at the same time. Notice that, although the sequences of widths look similar, they are not identical. This is typical 2. Measuring Ring Widths. Each core is sanded down with a belt sander using medium-grit paper and then finished by hand with flourgrade-grit paper. The rings are then clearly visible and differentiated from each other with a result very much like that shown in Figure A2. The core is then mounted on a movable table below a microscope and the ring-widths measured individually from the innermost ring to the outermost. The widths are automatically recorded in a computer file as they are measured (see Fig A3).

3. Cross-matching and Dating the Samples. Because of the factors besides the local climate which may determine the annual widths of a tree's rings, no two sequences of ring widths from different oaks growing at the same time are exactly alike (Fig A4). Indeed, the sequences may not be exactly alike even when the trees are growing near to each other. Consequently, in the Laboratory we do not attempt to match two sequences of ring widths by eye, or graphically, or by any other subjective method. Instead, it is done objectively (ie statistically) on a computer by a process called cross-matching. The output from the computer tells us the extent of correlation between two sample sequences of widths or, if we are dating, between a sample sequence of widths and the master, at each relative position of one to the other (offsets). The extent of the correlation at an offset is determined by the *t*-value (defined in almost any introductory book on statistics). That offset with the maximum t-value among the t-values at all the offsets will be the best candidate for dating one sequence relative to the other. If one of these is a master chronology, then this will date the other. Experiments carried out in the past with sequences from oaks of known date suggest that a t-value of at least 4.5, and preferably at least 5.0, is usually adequate for the dating to be accepted with reasonable confidence (Laxton and Litton 1988; Laxton et al 1988; Howard et al 1984–1995).

This is illustrated in Figure A5 with timbers from one of the roofs of Lincoln Cathedral. Here four sequences of ring widths, LIN-CO4, 05, 08, and 45, have been cross-matched with each other. The ring widths themselves have been omitted in the bar diagram, as is usual, but the offsets at which they best cross-match each other are shown; eg the sequence of ring widths of CO8 matches the sequence of ring widths of C45 best when it is at a position starting 20 rings after the first ring of C45, and similarly for the others. The actual *t*-values between the four at these offsets of best correlations are in the matrix. Thus at the offset of +20 rings, the *t*-value between C45 and C08 is 5.6 and is the maximum found between these two among all the positions of one sequence relative to the other.

It is standard practice in our Laboratory first to cross-match as many as possible of the ring-width sequences of the samples in a building and then to form an average from them. This average is called a site sequence of the building being dated and is illustrated in Figure A5. The fifth bar at the bottom is a site sequence for a roof at Lincoln Cathedral and is constructed from the matching sequences of the four timbers. The site sequence width for each year is the average of the widths in each of the sample sequences which has a width for that year. Thus in Fig A5 if the widths shown are 0.8mm for C45, 0.2mm for C08, 0.7mm for C05, and 0.3mm for C04, then the

corresponding width of the site sequence is the average of these, 0.55mm. The actual sequence of widths of this site sequence is stored on the computer. The reason for creating site sequences is that it is usually easier to date an average sequence of ring widths with a master sequence than it is to date the individual component sample sequences separately.

The straightforward method of cross-matching several sample sequences with each other one at a time is called the 'maximal *t*-value' method. The actual method of cross-matching a group of sequences of ring-widths used in the Laboratory involves grouping and averaging the ring-width sequences and is called the 'Litton-Zainodin Grouping Procedure'. It is a modification of the straightforward method and was successfully developed and tested in the Laboratory and has been published (Litton and Zainodin 1991; Laxton *et al* 1988).

4. Estimating the Felling Date. As mentioned above, if the bark is present on a sample, then the date of its last ring is the date of the felling of its tree (or the last full year before felling, if it was felled in the first three months of the following calendar year, before any new growth had started, but this is not too important a consideration in most cases). The actual bark may not be present on a timber in a building, though the dendrochronologist who is sampling can often see from its surface that only the bark is missing. In these cases the date of the last ring is still the date of felling.

Quite often some, though not all, of the original outer rings are missing on a timber. The outer rings on an oak, called sapwood rings, are usually lighter than the inner rings, the heartwood, and so are relatively easy to identify. For example, sapwood can be seen in the corner of the rafter and at the outer end of the core in Figure A2, both indicated by arrows. More importantly for dendrochronology, the sapwood is relatively soft and so liable to insect attack and wear and tear. The builder, therefore, may remove some of the sapwood for precisely these reasons. Nevertheless, if at least some of the sapwood rings are left on a sample, we will know that not too many rings have been lost since felling so that the date of the last ring on the sample is only a few years before the date of the original last ring on the tree, and so to the date of felling.

Various estimates have been made and used for the average number of sapwood rings in mature oak trees (English Heritage 1998). A fairly conservative range is between 15 and 50 and that this holds for 95% of mature oaks. This means, of course, that in a small number of cases there could be fewer than 15 and more than 50 sapwood rings. For example, the core CRO-A06 has only 9 sapwood rings and some have obviously been lost over time – either they were removed originally by the carpenter and/or they rotted away in the building and/or they were lost in the coring. It is not known exactly how many sapwood rings are missing, but using the above range the Laboratory would estimate between a minimum of 6 (=15-9) and a maximum of 41 (=50-9). If the last ring of CRO-A06 has been dated to 1500, say, then the estimated felling-date range for the tree from which it came originally would be between 1506 and 1541. The Laboratory uses this estimate for sapwood in areas of England where it has no prior information. It also uses it when dealing with samples with very many rings, about 120 to the last heartwood ring. But in other areas of England where the Laboratory has accumulated a number of samples with complete sapwood, that is, no sapwood lost since felling, other estimates in place of the conservative range of 15 to 50 are used. In the East Midlands (Laxton *et al* 2001) and the east to the south down to Kent (Pearson 1995) where it has sampled extensively in the past, the Laboratory uses the shorter estimate of 15 to 35 sapwood rings in 95% of mature oaks growing in these parts. Since the sample CRO-A06 comes from a house in Cropwell Bishop in the East Midlands, a better estimate of sapwood rings lost since felling is between a minimum of 6 (=15-9) and 26 (=35-9) and the felling would be estimated to have taken place between 1506 and 1526, a shorter period than before. Oak boards quite often come from the Baltic region and in these cases the 95% confidence limits for sapwood are 9 to 36 (Howard *et al* 1992, 56).

Even more precise estimates of the felling date and range can often be obtained using knowledge of a particular case and information gathered at the time of sampling. For example, at the time of sampling the dendrochronologist may have noted that the timber from which the core of Figure A2 was taken still had complete sapwood but that some of the soft sapwood rings were lost in coring. By measuring into the timber the depth of sapwood lost, say 20mm, a reasonable estimate can be made of the number of sapwood rings lost, say 12 to 15 rings in this case. By adding on 12 to 15 years to the date of the last ring on the sample a good tight estimate for the range of the felling date can be obtained, which is often better than the 15 to 35 years later we would have estimated without this observation. In the example, the felling is now estimated to have taken place between AD 1512 and 1515, which is much more precise than without this extra information.


Even if all the sapwood rings are missing on a sample, but none of the heartwood rings are, then an estimate of the felling-date range is possible by adding on the full compliment of, say, 15 to 35 years to the date of the last heartwood ring (called the heartwood/ sapwood boundary or transition ring and denoted H/S). Fortunately it is often easy for a trained dendrochronologist to identify this boundary on a timber. If a timber does not have its heartwood/sapwood boundary, then only a *post quem* date for felling is possible.

**5. Estimating the Date of Construction.** There is a considerable body of evidence collected by dendrochronologists over the years that oak timbers used in buildings were not seasoned in medieval or early modern times (English Heritage 1998; Miles 1997, 50–5). Hence, provided that all the samples in a building have estimated felling-date ranges broadly in agreement with each other, so that they appear to have been felled as a group, then this should give an accurate estimate of the period when the structure was built, or soon after (Laxton *et al* 2001, fig 8; 34–5, where 'associated groups of fellings' are discussed in detail). However, if there is any evidence of storage before use, or if there is evidence the oak came from abroad (eg Baltic boards), then some allowance has to be made for this.

6. Master Chronological Sequences. Ultimately, to date a sequence of ring widths, or a site sequence, we need a master sequence of dated ring widths with which to cross-match it, a Master Chronology. To construct such a sequence we have to start with a sequence of widths whose dates are known and this means beginning with a sequence from an oak tree whose date of felling is known. In Figure A6 such a sequence is SHE-T, which came from a tree in Sherwood Forest which was blown down in a recent gale. After this other sequences which cross-match with it are added and gradually the sequence is 'pushed back in time' as far as the age of samples will allow. This process is illustrated in Figure A6. We have a master chronological sequence of widths for Nottinghamshire and East Midlands oak for each year from AD 882 to 1981. It is described in great detail in Laxton and Litton (1988), but the components it contains are shown here in the form of a bar diagram. As can be seen, it is well replicated in that for each year in this period there are several sample sequences having widths for that year. The master is the average of these. This master can now be used to date oak from this area and from the surrounding areas where the climate is very similar to that in the East Midlands. The Laboratory has also constructed a master for Kent (Laxton and Litton 1989). The method the Laboratory uses to construct a master sequence, such as the East Midlands and Kent, is completely objective and uses the Litton-Zainodin grouping procedure (Laxton et al 1988). Other laboratories and individuals have constructed masters for other areas and have made them available. As well as these masters, local (dated) site chronologies can be used to date other buildings from nearby. The Laboratory has hundreds of these site sequences from many parts of England and Wales covering many short periods.


**Ring Width Indices.** Tree-ring dating can be done by cross-matching the ring 7. widths themselves, as described above. However, it is advantageous to modify the widths first. Because different trees grow at different rates and because a young oak grows in a different way from an older oak, irrespective of the climate, the widths are first standardized before any matching between them is attempted. These standard widths are known as ring-width indices and were first used in dendrochronology by Baillie and Pilcher (1973). The exact form they take is explained in this paper and in the appendix of Laxton and Litton (1988) and is illustrated in the graphs in Figure A7. Here ring-widths are plotted vertically, one for each year of growth. In the upper sequence of (a), the generally large early growth after 1810 is very apparent as is the smaller later growth from about 1900 onwards when the tree is maturing. A similar phenomenon can be observed in the lower sequence of (a) starting in 1835. In both the widths are also changing rapidly from year to year. The peaks are the wide rings and the troughs are the narrow rings corresponding to good and poor growing seasons, respectively. The two corresponding sequence of Baillie-Pilcher indices are plotted in (b) where the differences in the immature and mature growths have been removed and only the rapidly changing peaks and troughs remain, that are associated with the common climatic signal. This makes cross-matching easier.


t-value/offset Matrix




## Figure A5: Cross-matching of four sequences from a Lincoln Cathedral roof and the formation of a site sequence from them

The bar diagram represents these sequences without the rings themselves. The length of the bar is proportional to the number of rings in the sequence. Here the four sequences are set at relative positions (offsets) to each other at which they have maximum correlation as measured by the *t*-values. The *t*-value/offset matrix contains the maximum *t*-values below the diagonal and the offsets above it. Thus, the maximum *t*-value between C08 and C45 occurs at the offset of +20 rings and the *t*-value is then 5.6. The site sequence is composed of the average of the corresponding widths, as illustrated with one width







## Figure A7 (a): The raw ring-widths of two samples, THO-A01 and THO-B05, whose felling dates are known

Here the ring widths are plotted vertically, one for each year, so that peaks represent wide rings and troughs narrow ones. Notice the growth-trends in each; on average the earlier rings of the young tree are wider than the later ones of the older tree in both sequences

#### Figure A7 (b): The Baillie-Pilcher indices of the above widths

The growth trends have been removed completely

#### References

Baillie, M G L, and Pilcher, J R, 1973 A simple cross-dating program for tree-ring research, *Tree-Ring Bull*, **33**, 7–14

English Heritage, 1998 Dendrochronology: Guidelines on Producing and Interpreting Dendrochronological Dates, London

Hillam, J, Morgan, R A, and Tyers, I, 1987 Sapwood estimates and the dating of short ring sequences, *Applications of tree-ring studies*, BAR Int Ser, **3**, 165–85

Howard, R E, Laxton, R R, Litton, C D, and Simpson, W G, 1984–95 Nottingham University Tree-Ring Dating Laboratory results, *Vernacular Architect*, **15–26** 

Hughes, M K, Milson, S J, and Legett, P A, 1981 Sapwood estimates in the interpretation of tree-ring dates, *J Archaeol Sci*, **8**, 381–90

Laxon, R R, Litton, C D, and Zainodin, H J, 1988 An objective method for forming a master ring-width sequence, *P A C T*, **22**, 25–35

Laxton, R R, and Litton, C D, 1988 An East Midlands Master Chronology and its use for dating vernacular buildings, University of Nottingham, Department of Archaeology Publication, Monograph Series III

Laxton, R R, and Litton, C D, 1989 Construction of a Kent master dendrochronological sequence for oak, AD 1158 to 1540, *Medieval Archaeol*, **33**, 90–8

Laxton, R R, Litton, C D, and Howard, R E, 2001 *Timber: Dendrochronology of Roof Timbers at Lincoln Cathedral*, Engl Heritage Res Trans, 7

Litton, C D, and Zainodin, H J, 1991 Statistical models of dendrochronology, *J Archaeol Sci*, **18**, 29–40

Miles, D W H, 1997 The interpretation, presentation and use of tree-ring dates, Vernacular Architect, 28, 40–56

Pearson, S, 1995 The Medieval Houses of Kent, an Historical Analysis, London

Rackham, O, 1976 Trees and Woodland in the British Landscape, London



#### ENGLISH HERITAGE RESEARCH DEPARTMENT

English Heritage undertakes and commissions research into the historic environment, and the issues that affect its condition and survival, in order to provide the understanding necessary for informed policy and decision making, for sustainable management, and to promote the widest access, appreciation and enjoyment of our heritage.

The Research Department provides English Heritage with this capacity in the fields of buildings history, archaeology, and landscape history. It brings together seven teams with complementary investigative and analytical skills to provide integrated research expertise across the range of the historic environment. These are:

- \* Aerial Survey and Investigation
- \* Archaeological Projects (excavation)
- \* Archaeological Science
- \* Archaeological Survey and Investigation (landscape analysis)
- \* Architectural Investigation
- Imaging, Graphics and Survey (including measured and metric survey, and photography)
- \* Survey of London

The Research Department undertakes a wide range of investigative and analytical projects, and provides quality assurance and management support for externally-commissioned research. We aim for innovative work of the highest quality which will set agendas and standards for the historic environment sector. In support of this, and to build capacity and promote best practice in the sector, we also publish guidance and provide advice and training. We support outreach and education activities and build these in to our projects and programmes wherever possible.

We make the results of our work available through the Research Department Report Series, and through journal publications and monographs. Our publication Research News, which appears three times a year, aims to keep our partners within and outside English Heritage up-to-date with our projects and activities. A full list of Research Department Reports, with abstracts and information on how to obtain copies, may be found on www.english-heritage. org.uk/researchreports

For further information visit www.english-heritage.org.uk

