Some Notes on Lepidoptera, 1934. By H. C. Hayward, M.A., F.R.E.S.

I had hoped this year to pay more attention than previously to the lepidoptera of some of the Derbyshire Moors, to which I had been given access through the kindness of the Duke of Devonshire, but unfortunately, after a preliminary visit in April, I was unable to avail myself of the permission again, owing to ill-health. On this visit hibernated specimens of *Peronea mixtana* were present in some numbers but *Philodone prodromana* did not appear and the only other species observed was *Xanthorhoe multistrigaria*, of which a female was obtained which however refused to lay.

In the hopes of obtaining a few more melanics of Eupithecia trisignaria of which the two specimens bred last year and one previously are, so far as I know, the only recorded examples, a large number of larvae had been collected last autumn, but the results were disappointing, as out of 150 pupae none but normal forms were bred. My previous conviction of the comparative immunity of this species from parasites was, however, shaken, for a large number of parasites of at least two species were reared. In April larvae and pupae of Argyroploce nigricostana, a species I had frequently sought in vain in previous years, were found freely in a few spots in stems of Stachys sylvatica. As I was unable to do any active collecting after April, I fear I have no notes to record, except the very remarkable occurrence in Repton Shrubs in July of Argynnis paphia, of which one specimen was captured and three more seen by a boy in Repton School. The most recent record of this species in the county was of a single specimen near Chesterfield

in 1906, though it was recorded as not uncommon in Repton Shrubs and at Melbourne previously to 1880. is pleasant to find that it still lingers in the county though it is remarkable that so striking an insect should have escaped notice through so many years. As however this is the last occasion on which I shall contribute annual notes on Lepidoptera to this Journal, it has been suggested to me that it might be of interest to put together in a single article all the additional records for the county since the publication, thirty years ago, of the list compiled for the Victoria County History, by the Rev. F. C. R. Jourdain. Most of these are included in a list published by the Derbyshire Entomological Society in 1926, but that list is less well known and less readily accessible than the V.C.H. The order and nomenclature adopted is that of Meyricks Revised Handbook and I have added the year and locality in which each species was first observed. Most of the new records from the north of the county have been contributed by Mr. H. W. Daltry of Madeley and to these his initials are appended. In the case of a few other species the observer's name has been added: the remaining records are my own.

CARADRININA.

Hadena leucostigma Hb. Repton, 1912.

Hadena bicoloria Vill. Willington, 1905.

Euxoa puta Hb. Repton, 1931.

Agrotis lucernea L. Dovedale, 1931. H.W.D.

Polia lichenea Hb. Mill Dale, 1922. J. Hill.

Monima opima Hb. Willington, 1909.

Melanchria serena F. Repton, 1914.

Plusia moneta F. Repton, 1906.

NOTODONTINA.

Leucopthalmia annulata Sch. Repton, 1915.

Trichopteryx carpinata Bkh. *Chesterfield*, 1889. S. Hooke.

Chlorochlystis coronata Hb. Coxbench, 1916. G. H.

Sale.

Eupithecia pusillata F. Repton, 1920.

Deileptenia abietaria Hb. Repton, 1911.

Polyploca ridens F. Chesterfield, 1909. S. Hooke.

Deilephila lineata F. Repton, 1931.

PAPILIONINA.

Limenitis sibilla L. Repton, 1910.

Hesperia malvae L. Willington, 1919.

PYRALIDINA.

Phycita spissicella F. Repton, 1916.

Ephestia cahiritella Z. Derby, 1922.

Ephestia Kühniella Z. Repton, 1920.

Acrobasis Zelleri Rag. Repton, 1919.

Crambus pinellus L. Repton, 1917.

Scoparia crataegella Hb. Willington, 1916.

Scoparia angustea Stph. Repton, 1926.

Oxyptilus heterodactylus Vill. Ticknall, 1928.

Marasmarcha microdactyla Hb. Dovedale, 1926.

H.W.D.

Alucita baliodactyla Z. Dovedale, 1926. H.W.D. TORTRICINA.

Phalonia smeathmanniana F. Staveley. L. Wyre.

Phalonia dubitana Hb. Willington, 1917.

Chlidonia subbaumanniana Wilk. Via Gellia, 1917.

Chlidonia baumanniana Schiff. Via Gellia, 1917.

Eulia politana Haw. Holymoor, 1922.

Cnephasia pascuana Hb. Repton, 1916.

Peronea aspersana Hb. Repton, 1918.

Peronea mixtana Hb. Matlock, 1926.

Peronea hastiana L. Repton. R. E. E. Frampton.

Peronea caledoniana Stph. Matlock, 1931.

Spilonota lariciana Z. Repton, 1928.

Evetria posticana Zett. Repton, 1916.

Ancylis siculana Hb. Dovedale, 1926. H.W.D.

Eucosma rubiginosana HS. Repton, 1917.

Eucosma expallidana Haw. Repton, 1933.

Eucosma turbidana Tr. Dovedale, 1927. H.W.D. Eucosma cinereana Haw. Repton, 1916.
Eucosma semifuscana Stph. Repton, 1916.
Argyroploce bifasciana Haw. Repton, 1917.
Hemimene politana Hb. Willington, 1917.
Hemimene aplinana Tr. Dovedale, 1931. H.W.D. Hemimene tanaceti Stt. Willington, 1917.
Pammene ochsenheimeriana Z. Repton, 1922.
Pammene splendidulana Gn. Repton, 1928.
Pammene fimbriana Haw. Repton, 1920.
Pammene populana F. Repton, 1924.
Laspeyresia coniferana Rtzb. Repton, 1925.
Laspeyresia cosmophorana Tr. Matlock, 1933.
Tineina.

Sitotroga cerealella Ol. Repton, 1921. Stenolechia gemmella L. Repton, 1916. Exoteleia dodecella L. Repton, 1922. Telphusa scalella Sc. Repton, 1917. Telphusa sequax Haw. Dovedale, 1927. H.W.D. Gelechia similis St. Dovedale, 1929. H.W.D. Gelechia politella Stt. Dovedale, 1927. H.W.D. Gelechia diffinis Haw. Willington, 1919. Gelechia longicornis Curt. Holymoor, 1922. Gelechia rhombella Schiff. Repton, 1930. Gelechia nigra Haw. Repton, 1916. Ysolophus semicostellus Hb. Dovedale, 1927. H.W.D. Schiffermuelleria augustella Hb. Willington, 1918. Borkhausenia subaquilea St. Matlock, 1933. Cheimophila salicella Hb. Willington, 1916. Pleurota bicostella Cl. Breadsall, 1917. Depressaria zephyrella Hb. Repton, 1916. Depressaria conterminella Z. Repton, 1918. Aegeria formiciformis Esp. Repton, 1909.

¹ This is the species recently differentiated from H. flavidorsana, Knaggs, to which Mr. Daltry's specimen was first ascribed. Meyrick's nomenclature is here incorrect and confused. He calls H. politana *quaestionana*, a synonmy of alpinana.

Heliozele stannella F.R. Repton, 1916. Pancalia leuwenhoekella L. Dovedale, 1927. H.W.D. Schreckensteinia festaliella Hb. Repton, 1916. Perittia obscuripunctella Stt. Willington, 1920. Stephensia brunnichiella L. Tichnall, 1928. Elachista gleichenella F. Repton, 1920. Elachista apicipunctella Stt. Repton, 1916. Elachista luticomella Z. Dovedale, 1927. H.W.D. Elachista triatomea Haw. Dovedale, 1927. H.W.D. Cedestis farinatella Dup. Repton, 1916. Cedestis gysselinella Dup. Repton, 1916. Argyresthia atmoriella Banks. Repton, 1916. Argyresthia pygmaeella Haw. Repton, 1917. Argyresthia mendica Haw. Repton, 1916. Argyresthia semifusca Haw. Repton, 1916. Argyresthia conjugella Z. Repton, 1916. Swammerdamia combinella Hb. Willington, 1918. Prays curtisellus Don. Repton, 1916. Hyponomeuta stannella Thnb. Dovedale, 1926. H.W.D.

(The first record of this species in Britain). Ethmia funerella F. Via Gellia, 1917. Coleophora lutipennella Z. Repton, 1916. Coleophora palliatella Zk. Repton, 1922. Lithocolletis spinicolella Stt. Willington, 1919. Lithocolletis froelichiella Z. Willington, 1916. Lithocelletis stettinensis Nic. Willington, 1919. Ornix scoticella Stt. Repton, 1920. Ornix betulae Stt. Repton, 1916. Gracilaria hemidactylella F. Repton, 1922. Cataplectica fulviguttella Z. Repton, 1932. Cerostoma nemorella L. Repton, 1916. Tineola biseliella Hm. Repton, 1916. Ochsenheimeria bisontella Z. Dovedale, 1927. H.W.D. Fumea intermediella Brd. Repton, 1916. Incurvaria rubiella Bjerk. Repton, 1916.

Adela cuprella Thnb. Willington, 1918. NEPTICULINA.

Nepticula pulverosella Stt. *Repton*, 1916. MICROPTERYGINA.

Eriocrania semipurpurella Stph. Willington, 1917. Eriocrania purpurella Haw. Repton, 1916. Mnemonica unimaculella Zett. Repton, 1916.