



ARCHAEOLOGICAL
SERVICES
WYAS

**Land south of The Hollies
Burgh-le-Marsh
Lincolnshire**

Geophysical Survey

Report no. 2499

July 2013

Client: CgMs Consulting



Land south of The Hollies
Burgh-le-Marsh
Lincolnshire

Geophysical Survey

Summary

A geophysical (magnetometer) survey, covering approximately 3.9 hectares, was carried out at the The Hollies, Burgh-le-Marsh, in advance of a proposed development. The survey has demonstrated that the site is located across the geological boundary between superficial deposits of till and tidal flat deposits, and that, prior to drainage, much of the site is likely to have formed part of a wetland environment. No anomalies of archaeological potential have been identified by the survey and, therefore, the archaeological potential of the site is considered to be low.



ARCHAEOLOGICAL
SERVICES
WYAS

Report Information

Client: CgMs Consulting
Address: Burlington House, Lypiatt Road, Cheltenham,
Gloucestershire, GL50 2SY
Report Type: Geophysical Survey
Location: The Hollies, Burgh-le-Marsh
County: Lincolnshire
Grid Reference: TF 4970 6290
Period(s) of activity: -
Report Number: 2499
Project Number: 4087
Site Code: LSH13
OASIS ID: archaeol11-156501
Planning Application No.: -
Museum Accession No.: n/a
Date of fieldwork: July 2013
Date of report: July 2013
Project Management: Sam Harrison BSc MSc AIfA
Fieldwork: James Lawton BSc MSc PIfA
Kieron Kinninmont BSc
Report: David Harrison BA MSc MIfA
Illustrations: Christopher Sykes BA MSc
Photography: Site Staff

Authorisation for
distribution: _____



© Archaeological Services WYAS 2013
PO Box 30, Nephshaw Lane South, Morley, Leeds LS27 0UG
Telephone: 0113 383 7500.
Email: admin@aswyas.com



Contents

Report information	ii
Contents.....	iii
List of Figures	iv
List of Plates	iv
1 Introduction	1
Site location, topography and land-use.....	1
Geology and soils.....	1
2 Archaeological background	1
3 Aims, Methodology and Presentation	2
4 Results and Discussion	3
5 Conclusions.....	4

Figures

Plates

Appendices

Appendix 1: Magnetic survey: technical information

Appendix 2: Survey location information

Appendix 3: Geophysical archive

Bibliography

List of Figures

- 1 Site location (1:50000)
- 2 Site location showing superficial geology (1:625000) and greyscale magnetometer data (1:5000)
- 3 Processed greyscale magnetometer data (1:1000)
- 4 XY trace plot of minimally processed magnetometer data (1:1000)
- 5 Interpretation of magnetometer data (1:1000)

List of Plates

- Plate 1 General view of survey area, looking south-west
Plate 2 General view of survey area, looking south

1 Introduction

Archaeological Services WYAS was commissioned by William Bedford of CgMs Consulting to carry out a programme of non-intrusive geophysical (magnetometer) survey on land south of The Hollies, Burgh-le-Marsh, Lincolnshire (see Fig. 1), in advance of the determination of a planning application for a proposed development. The work was undertaken in accordance with guidance contained within the National Planning Policy Framework (2012) and with a Project Design submitted to and approved by the client (Harrison, 2013). The survey was carried out on July 9th 2013.

Site location, topography and land-use

The site is located approximately 1km to the south of Burgh-le-Marsh (see Fig. 1) and 5km west of Skegness. The proposed development area (PDA) is contained within a single field to the south-west of The Hollies Country Cottages, off High Lane. It is bound to the north by landscaped ground incorporating a pond, wooded area and car park, and by open pasture on all other sides (see Fig. 2). The site lies on a slight south-facing gradient at 2m above Ordnance Datum (aOD) within the south of the PDA and 3m aOD in the north. At the time of survey the field was under a recently harvested bean crop (see plates).

Geology and soils

The underlying bedrock comprises Claxby Ironstone Formation, Tealby Formation and Roach Formation overlain by till in the north of the survey area and tidal flat deposits in the south (British Geological Survey 2013). The soils in this area are classified in Holderness association, characterised as slowly permeable seasonally waterlogged fine loams (Soil Survey of England and Wales 1983).

2 Archaeological background

Research undertaken as background for the Environmental Statement (ECUS 2012) for a solar park scheme to the north-west of The Hollies identified that there are numerous archaeological sites known in the wider landscape and, in the immediate vicinity, several cropmark sites indicative of medieval settlement. The Archaeology and Cultural Heritage chapter of the Environmental Statement concluded that the 'archaeological potential for the proposed scheme is considered to be moderate'. Subsequent geophysical survey geophysical (magnetometer) survey carried out prior to trial trenching (Webb 2012) confirmed the presence of a square enclosure with internal divisions and numerous discrete features (see Fig 2). In addition, a plethora of linear and curvilinear anomalies in the south-eastern corner of the site located a series of enclosures, previously unknown, thought to be indicative of an area of settlement of presumed Roman/early post-Roman date. The trial trench evaluation (Richardson *et al* 2012) based on the previous geophysics identified features consistent with an extensive Roman settlement, with later Iron Age origins, to the north of The Hollies as

indicated by the geophysical survey. An enclosure of medieval date has also been confirmed. All of the trenches located in apparently 'blank' areas were found to be devoid of archaeological remains confirming that the geophysical survey provided an accurate assessment of the extent of the archaeological resource on this site.

3 Aims, Methodology and Presentation

The general objective of the geophysical survey was to provide information about the presence/absence, character, and extent of any archaeological remains identified within the specific area to be impacted by the proposed development and to help inform further strategies should they be required.

In order to achieve these aims detailed (recorded) magnetometer survey was carried out over the PDA, an area of 3.9 hectares.

The survey area was set-out with a Trimble 5600 Total Station Theodolite and the site grid tied into permanent landscape features. The grid was then superimposed onto digital Ordnance Survey mapping supplied by the client. Temporary reference objects (wooden survey marker stakes) were established and left in place following completion of the fieldwork for accurate geo-referencing. The locations of the temporary reference objects are shown on Figure 2 and their Ordnance Survey co-ordinates tabulated in Appendix 2.

Magnetometer survey

Bartington Grad601 magnetic gradiometers were used during the survey taking readings at 0.25m intervals on zig-zag traverses 1m apart within 30m by 30m grids so that 3600 readings were recorded in each grid. These readings were stored in the memory of the instrument and later downloaded to computer for processing and interpretation. Geoplot 3 (Geoscan Research) software was used to process and present the data. Further details are given in Appendix 1.

Reporting

A general site location plan, incorporating the 1:50000 Ordnance Survey map is shown in Figure 1. A large scale (1:5000) site location plan showing the greyscale magnetometer data and the approximate location of the superficial geology boundary (1:625000) is shown in Figure 2. Detailed plots of the data are presented in greyscale, XY trace plot and interpretation formats in Figures 3, 4 and 5 at a scale of 1:1000.

Further technical information on the equipment used, data processing and survey methodologies are given in Appendix 1 and Appendix 2. Appendix 3 describes the composition and location of the site archive.

The geophysical survey methodology, report and any recommendations comply with guidelines outlined by English Heritage (David *et al.* 2008) and by the Institute for Archaeologists (IfA 2010). All figures reproduced from Ordnance Survey mapping are with the permission of the controller of Her Majesty's Stationery Office (© Crown copyright).

The figures in this report have been produced following analysis of the data in 'raw' and processed formats and over a range of different display levels. All figures are presented to most suitably display and interpret the data from this site based on the experience and knowledge of Archaeological Services staff.

4 Results and Discussion

The geophysical (magnetometer) survey has identified a variable background response, particularly within the southern half of the PDA. The anomalies identified by the survey fall into three different categories according to their origin and these are discussed below.

Ferrous anomalies

Ferrous anomalies, either as individual 'spikes' or more extensive areas of magnetic disturbance, are typically caused by ferrous (magnetic) debris, either on the ground surface or mixed in with the plough-soil. Little importance is normally given to such anomalies, unless there is any supporting evidence for an archaeological interpretation, as ferrous debris is common on rural sites, often being present as a consequence of manuring or tipping/infilling. On this site there is no clustering to these anomalies and they are therefore assumed to be due to the random distribution of ferrous debris.

An area of magnetic disturbance at the north-eastern corner of the survey area is caused by the proximity of a pylon and a gate within the adjacent field boundary (see Figs 3, 4 and 5).

Geological anomalies

Amorphous areas of magnetic enhancement dominate the dataset, although a less variable background is clearly discernable within the north of the PDA. This difference in magnetic background corresponds to the superficial geological boundary between till, in the north of the PDA, and tidal flat deposits to the south. The anomalies are interpreted as being due to areas of alluvium and magnetically enhanced silts being deposited on the lower-lying part of the site following episodes of flooding. It is worth considering here that the magnetometer detects anomalies up to a depth of approximately 1m and that the continuous build up of alluvial deposits may result in discrete anomalies of archaeological potential remaining beyond detection. The broad, linear anomaly, **A**, within the north of the PDA is likely to be caused by a former water channel.

Agricultural anomalies

A dipolar linear anomaly, **B**, is visible crossing the PDA on a north-west/south-east alignment. This anomaly corresponds to a former field boundary depicted on the first edition Ordnance Survey map of 1889, and whilst it may be of local historical interest, it is unlikely to be of any archaeological interest. Towards the south of the site, a discrete linear trend, **C**, is likely to locate the route of a field drain.

5 Conclusions

The survey has identified that the PDA lies on the superficial geological boundary between tidal flats in the south and till to the north. Marginal lands surrounding former wetland environments are likely to have been utilised in the past, particularly in prehistory and, as discussed, extensive Romano-British and Roman settlement activity is recorded on the higher ground to the immediate north of the PDA. However, no anomalies of archaeological potential have been identified by the geophysical survey, and therefore, on this basis, the PDA is assessed as having a low archaeological potential.

The results and subsequent interpretation of data from geophysical surveys should not be treated as an absolute representation of the underlying archaeological and non-archaeological remains. Confirmation of the presence or absence of archaeological remains can only be achieved by direct investigation of sub-surface deposits.





Fig. 1. Site location




Fig. 2. Site location showing superficial geology (1:625000) and greyscale magnetometer data (1:5000 @ A3)

0 50m




Fig. 3. Processed greyscale magnetometer data (1:1000 @ A3)





Fig. 4. XY trace plot of minimally processed magnetometer data (1:1000 @ A3)



Reproduced from the Ordnance Survey mapping with the permission of the Controller of Her Majesty's Stationery Office. © Crown Copyright. Unauthorised reproduction infringes Crown copyright and may lead to prosecution or civil proceedings. Digital data supplied by the client, CgMs, 2013.

© ASWYAS 2013. Archaeological Services W Y A S PO Box 30, Nephaw Lane South, Morley, LS27 0UG Tel: 0113 383 7500 Fax: 0113 383 7501

Fig. 5. Interpretation of magnetometer data (1:1000 @ A4)

0 25m



Plate 1. General view of survey area, looking south-west



Plate 2. General view of survey area, looking south

Appendix 1: Magnetic survey - technical information

Magnetic Susceptibility and Soil Magnetism

Iron makes up about 6% of the Earth's crust and is mostly present in soils and rocks as minerals such as maghaemite and haemetite. These minerals have a weak, measurable magnetic property termed magnetic susceptibility. Human activities can redistribute these minerals and change (enhance) others into more magnetic forms so that by measuring the magnetic susceptibility of the topsoil, areas where human occupation or settlement has occurred can be identified by virtue of the attendant increase (enhancement) in magnetic susceptibility. If the enhanced material subsequently comes to fill features, such as ditches or pits, localised isolated and linear magnetic anomalies can result whose presence can be detected by a magnetometer (fluxgate gradiometer).

In general, it is the contrast between the magnetic susceptibility of deposits filling cut features, such as ditches or pits, and the magnetic susceptibility of topsoils, subsoils and rocks into which these features have been cut, which causes the most recognisable responses. This is primarily because there is a tendency for magnetic ferrous compounds to become concentrated in the topsoil, thereby making it more magnetic than the subsoil or the bedrock. Linear features cut into the subsoil or geology, such as ditches, that have been silted up or have been backfilled with topsoil will therefore usually produce a positive magnetic response relative to the background soil levels. Discrete feature, such as pits, can also be detected. The magnetic susceptibility of a soil can also be enhanced by the application of heat and the fermentation and bacterial effects associated with rubbish decomposition. The area of enhancement is usually quite large, mainly due to the tendency of discard areas to extend beyond the limit of the occupation site itself, and spreading by the plough. An advantage of magnetic susceptibility over magnetometry is that a certain amount of occupational activity will cause the same proportional change in susceptibility, however weakly magnetic is the soil, and so does not depend on the magnetic contrast between the topsoil and deeper layers. Susceptibility survey is therefore able to detect areas of occupation even in the absence of cut features. On the other hand susceptibility survey is more vulnerable to the masking effects of layers of colluvium and alluvium as the technique, using the Bartington system, can generally only measure variation in the first 0.15m of ploughsoil.

Types of Magnetic Anomaly

In the majority of instances anomalies are termed 'positive'. This means that they have a positive magnetic value relative to the magnetic background on any given site. However some features can manifest themselves as 'negative' anomalies that, conversely, means that the response is negative relative to the mean magnetic background.

Where it is not possible to give a probable cause of an observed anomaly a '?' is appended.

It should be noted that anomalies interpreted as modern in origin might be caused by features that are present in the topsoil or upper layers of the subsoil. Removal of soil to an archaeological or natural layer can therefore remove the feature causing the anomaly.

The types of response mentioned above can be divided into five main categories that are used in the graphical interpretation of the magnetic data:

Isolated dipolar anomalies (iron spikes)

These responses are typically caused by ferrous material either on the surface or in the topsoil. They cause a rapid variation in the magnetic response giving a characteristic 'spiky' trace. Although ferrous archaeological artefacts could produce this type of response, unless there is supporting evidence for an archaeological interpretation, little emphasis is normally given to such anomalies, as modern ferrous objects are common on rural sites, often being present as a consequence of manuring.

Areas of magnetic disturbance

These responses can have several causes often being associated with burnt material, such as slag waste or brick rubble or other strongly magnetised/fired material. Ferrous structures such as pylons, mesh or barbed wire fencing and buried pipes can also cause the same disturbed response. A modern origin is usually assumed unless there is other supporting information.

Linear trend

This is usually a weak or broad linear anomaly of unknown cause or date. These anomalies are often caused by agricultural activity, either ploughing or land drains being a common cause.

Areas of magnetic enhancement/positive isolated anomalies

Areas of enhanced response are characterised by a general increase in the magnetic background over a localised area whilst discrete anomalies are manifest by an increased response (sometimes only visible on an XY trace plot) on two or three successive traverses. In neither instance is there the intense dipolar response characteristic exhibited by an area of magnetic disturbance or of an 'iron spike' anomaly (see above). These anomalies can be caused by infilled discrete archaeological features such as pits or post-holes or by kilns. They can also be caused by pedological variations or by natural infilled features on certain geologies. Ferrous material in the subsoil can also give a similar response. It can often therefore be very difficult to establish an anthropogenic origin without intrusive investigation or other supporting information.

Linear and curvilinear anomalies

Such anomalies have a variety of origins. They may be caused by agricultural practice (recent ploughing trends, earlier ridge and furrow regimes or land drains), natural geomorphological features such as palaeochannels or by infilled archaeological ditches.

Methodology: Magnetic Susceptibility Survey

There are two methods of measuring the magnetic susceptibility of a soil sample. The first involves the measurement of a given volume of soil, which will include any air and moisture that lies within the sample, and is termed volume specific susceptibility. This method results in a bulk value that is not necessarily fully representative of the constituent components of the sample. For field surveys a Bartington MS2 meter with MS2D field loop is used due to its speed and simplicity. The second technique overcomes this potential problem by taking into account both the volume and mass of a sample and is termed mass specific susceptibility. However, mass specific readings cannot be taken in the field where the bulk properties of a soil are usually unknown and so volume specific readings must be taken. Whilst these values are not fully representative they do allow general comparisons across a site and give a broad indication of susceptibility changes. This is usually enough to assess the susceptibility of a site and evaluate whether enhancement has occurred.

Methodology: Gradiometer Survey

There are two main methods of using the fluxgate gradiometer for commercial evaluations. The first of these is referred to as *magnetic scanning* and requires the operator to visually identify anomalous responses on the instrument display panel whilst covering the site in widely spaced traverses, typically 10m apart. The instrument logger is not used and there is therefore no data collection. Once anomalous responses are identified they are marked in the field with bamboo canes and approximately located on a base plan. This method is usually employed as a means of selecting areas for detailed survey when only a percentage sample of the whole site is to be subject to detailed survey.

The disadvantages of magnetic scanning are that features that produce weak anomalies (less than 2nT) are unlikely to stand out from the magnetic background and so will be difficult to detect. The coarse sampling interval means that discrete features or linear features that are parallel or broadly oblique to the direction of traverse may not be detected. If linear features are suspected in a site then the traverse direction should be perpendicular (or as close as is possible within the physical constraints of the site) to the orientation of the suspected features. The possible drawbacks mentioned above mean that a 'negative' scanning result should be validated by sample detailed magnetic survey (see below).

The second method is referred to as *detailed survey* and employs the use of a sample trigger to automatically take readings at predetermined points, typically at 0.25m intervals, on zig-zag traverses 1m apart. These readings are stored in the memory of the instrument and are later dumped to computer for processing and interpretation. Detailed survey allows the visualisation of weaker anomalies that may not have been detected by magnetic scanning.

During this survey a Bartington Grad601 magnetic gradiometer was used taking readings on the 0.1nT range, at 0.25m intervals on zig-zag traverses 1m apart within 30m by 30m square

grids. The instrument was checked for electronic and mechanical drift at a common point and calibrated as necessary. The drift from zero was not logged.

Data Processing and Presentation

The detailed gradiometer data has been presented in this report in XY trace and greyscale formats. In the former format the data shown is 'raw' with no processing other than grid biasing having been done. The data in the greyscale images has been interpolated and selectively filtered to remove the effects of drift in instrument calibration and other artificial data constructs and to maximise the clarity and interpretability of the archaeological anomalies.

An XY plot presents the data logged on each traverse as a single line with each successive traverse incremented on the Y-axis to produce a 'stacked' plot. A hidden line algorithm has been employed to block out lines behind major 'spikes' and the data has been clipped. The main advantage of this display option is that the full range of data can be viewed, dependent on the clip, so that the 'shape' of individual anomalies can be discerned and potentially archaeological anomalies differentiated from 'iron spikes'. Geoplot 3 software was used to create the XY trace plots.

Geoplot 3 software was used to interpolate the data so that 3600 readings were obtained for each 30m by 30m grid. The same program was used to produce the greyscale images. All greyscale plots are displayed using a linear incremental scale.

Appendix 2: Survey location information

The site grid was laid out using a Trimble VRS differential Global Positioning System (Trimble 5800 model). The accuracy of this equipment is better than 0.01m. The locations of the temporary reference points left on site are shown on Figure 2 and the Ordnance Survey grid co-ordinates tabulated below. The internal accuracy of these markers is better than 0.01m. The survey grids were then super-imposed onto a base map provided by the client to produce the displayed block locations. However, it should be noted that Ordnance Survey positional accuracy for digital map data has an error of 0.5m for urban and floodplain areas, 1.0m for rural areas and 2.5m for mountain and moorland areas. This potential error must be considered if co-ordinates are measured off hard copies of the mapping rather than using the digital co-ordinates.

Temporary reference objects were left on site (see Fig. 2). The Ordnance Survey reference points are listed below.

Station	Easting	Northing
A	549759.8305	362835.0905
B	549756.0672	362807.7616
C	549612.3108	362883.0385

Archaeological Services WYAS cannot accept responsibility for errors of fact or opinion resulting from data supplied by a third party. Archaeological Services WYAS cannot accept responsibility for errors of fact or opinion resulting from data supplied by a third party.

Archaeological Services WYAS cannot accept responsibility for errors of fact or opinion resulting from data supplied by a third party or for the removal of any of the survey reference points.

Appendix 3: Geophysical archive

The geophysical archive comprises:-

- an archive disk containing compressed (WinZip 8) files of the raw data, report text (Microsoft Word 2000), and graphics files (Adobe Illustrator CS2 and AutoCAD 2008) files; and
- a full copy of the report.

At present the archive is held by Archaeological Services WYAS although it is anticipated that it may eventually be lodged with the Archaeology Data Service (ADS). Brief details may also be forwarded for inclusion on the English Heritage Geophysical Survey Database after the contents of the report are deemed to be in the public domain (i.e. available for consultation in the Lincolnshire Historic Environment Record).

Bibliography

- British Geological Survey, 2013, <http://mapapps.bgs.ac.uk/geologyofbritain/home.html>
(Accessed: July 11th 2013)
- David, A., N. Linford, P. Linford and L. Martin, 2008, *Geophysical Survey in Archaeological Field Evaluation: Research and Professional Services Guidelines (2nd edition)* English Heritage
- ECUS, 2012, *The Hollies Solar Park. Environmental Report*. Unpublished Client Report No. 4433
- Harrison, S., 2013, *Land south of The Hollies, Burgh le Marsh, Lincolnshire; Geophysical Survey Project Design* Archaeological Services WYAS Unpublished document
- Institute for Archaeologists, 2010, *Standard and Guidance for archaeological geophysical survey*. Institute for Archaeologists
- Richardson, J., Webb, A. and Williams, D., 2012, *The Hollies Solar Park, Burgh-Le-Marsh, Lincolnshire: Trial Trench Evaluation* Archaeological Services WYAS Report no. 2477
- Soil Survey of England and Wales, 1983, Soils of Northern England Sheet 1
- Webb, A., 2012, *The Hollies Solar Park, Burgh-Le-Marsh, Lincolnshire: Geophysical Survey* Archaeological Services WYAS Report no. 2394