



ARCHAEOLOGICAL
SERVICES
WYAS

Skipton Flood Alleviation Scheme
Skipton
North Yorkshire

Geophysical Survey

Report no. 2561

December 2013

Client: Environment Agency



Skipton Flood Alleviation Scheme

North Yorkshire

Geophysical Survey

Summary

A geophysical (magnetometer) survey covering 4.5 hectares was carried out at Eller Beck and Waller Hill Beck, Skipton, in advance of ground works to facilitate the Skipton Flood Alleviation Scheme. Anomalies indicative of ridge and furrow cultivation have been identified at Waller Hill Beck, although the archaeological significance of these anomalies is thought to be limited. At Eller Beck, anomalies associated with a geological fault have been recorded. No anomalies of archaeological potential have been identified. Therefore, on the basis of the magnetic survey the archaeological potential of the two sites is considered to be low.



ARCHAEOLOGICAL
SERVICES
WYAS

Report Information

Client: Environment Agency
Address: NEAS. Lateral, 8 City Walk, Leeds LS11 9AT
Report Type: Geophysical Survey
Location: Eller Beck and Waller Hill Beck, Skipton
County: North Yorkshire
Grid Reference: Eller Beck: SD 9948 5505; Waller Hill Beck: SE 0060 5236
Period(s) of activity: Post Medieval?
Report Number: 2561
Project Number: 4152
Site Code: SKP13
OASIS ID: archaeol11- 168438
Planning Application No.:
Museum Accession No.: n/a
Date of fieldwork: November 2013
Date of report: December 2013
Project Management: Sam Harrison BSc MSc MifA
Fieldwork: Christopher Sykes BA MSc
Daniel Waterfall BA
Thomas Fildes BA
Report: David Harrison BA MSc MifA
Illustrations: David Harrison
Photography: Site Staff
Research: n/a

Authorisation for
distribution: -----



© Archaeological Services WYAS 2013
PO Box 30, Nepshaw Lane South, Morley, Leeds
LS27 0UG
Telephone: 0113 383 7500.
Email: admin@aswyas.com



Contents

Report information	ii
Contents.....	iii
List of Figures	iv
List of Plates	iv
1 Introduction	1
Site location, topography and land-use	1
Soils and Geology.....	1
2 Archaeological Background.....	1
3 Aims, Methodology and Presentation	2
4 Results and Discussion.....	3
5 Conclusions.....	3

Figures

Plates

Appendices

Appendix 1: Magnetic survey: technical information

Appendix 2: Survey location information

Appendix 3: Geophysical archive

Bibliography

List of Figures

- 1 Site location (1:50000)
- 2 Survey location showing greyscale magnetometer data (1:4000)
- 3 Processed greyscale magnetometer data; Eller Beck (1:1000)
- 4 XY trace plot of minimally processed magnetometer data; Eller Beck (1:1000)
- 5 Interpretation of magnetometer data; Eller Beck (1:1000)
- 6 Processed greyscale magnetometer data; Waller Hill Beck (1:1000)
- 7 XY trace plot of minimally processed magnetometer data; Waller Hill Beck (1:1000)
- 8 Interpretation of magnetometer data; Waller Hill Beck (1:1000)

List of Plates

- Plate 1 General view of Eller Beck survey area, looking north
- Plate 2 General view of Waller Hill Beck survey area, looking west
- Plate 3 General view of Waller Hill Beck survey area, looking north-east
- Plate 4 General view of Waller Hill Beck survey area, looking west

1 Introduction

Archaeological Services WYAS (ASWYAS) were commissioned by Emma Morrish of the Environment Agency to undertake a geophysical (magnetometer) survey of two sites to the north-west of Skipton. The survey was commissioned in advance of the construction of a flood storage bund at Eller Beck and both the construction a flood storage bund and excavation of a borrow pit at Waller Hill Beck to facilitate the Skipton Flood Alleviation Scheme. The work was undertaken in accordance with a Project Design (Harrison 2013), with guidance contained within the National Planning Policy Framework (2012) and in line with current best practice (David *et al* 2008). The survey was carried out on December 9th 2013 in order to provide additional information on the archaeological potential of the site.

Site location, topography and land-use

The Eller Beck site is located within a single field of pasture to the immediate north of the A65 and east of Skipton Golf Club's club house. A railway line borders the field to the north (see Fig 2; Plate 1). The site is centred at SD 9948 5505 and lies at approximately 137m above Ordnance Datum (aOD).

The Waller Hill Beck site is located to the south of the A65 and immediately north of Otley Road (A6069). It comprises three fields of pasture centred at SE 0060 5236 and laying at approximately 160m aOD (see Fig 2; Plate 2, Plate 3 and Plate 4).

Soils and geology

The underlying bedrock at the Eller Beck site comprises Clitheroe Limestone Formation – Mudstone overlain by superficial deposits of till. At the Waller Hill Beck site, the underlying bedrock comprises Clitheroe Limestone Formation – Mudstone, Chatburn Limestone Formation – Limestone and Embsay Limestone Member – Limestone overlain by till superficial deposits. A geological fault (see Fig. 2) is recorded on a north-west/south-east orientation passing through both sites (British Geological Survey 2013).

The soils in both areas are classified in the Brickfield 2 association, characterised as slowly permeable, seasonally waterlogged fine loams (Soil Survey of England and Wales 1983).

2 Archaeological Background

No archaeological remains are known from within the two sites or their immediate environs. An archaeological watching brief undertaken during the excavation of 38 geotechnical test pits, as part of this scheme, identified no archaeological remains (ASWYAS *In Prep.*). However, the wider landscape has produced evidence of occupation from the Mesolithic period onwards. Skipton itself was probably established in the early medieval period, becoming the location of a castle and associated settlement soon after the Norman conquest.

3 Aims, Methodology and Presentation

The main aim of the geophysical survey was to provide sufficient information to enable an assessment to be made of the impact of potential sub-surface archaeological remains and for further evaluation or mitigation proposals, if appropriate, to be recommended. To achieve this aim a magnetometer survey covering 1 hectare was undertaken at the Eller Beck site and a survey of 3.5 hectares was undertaken at the Waller Hill Beck site.

The general objectives of the geophysical survey were:

- to provide information about the nature and possible interpretation of any magnetic anomalies identified;
- to therefore determine the presence/absence and extent of any buried archaeological features; and
- to prepare a report summarising the results of the survey.

Magnetometer survey

The site grid was laid out using a Trimble VRS differential Global Positioning System (Trimble 5800 model). Bartington Grad601 magnetic gradiometers were used during the survey, taking readings at 0.25m intervals on zig-zag traverses 1m apart within 30m by 30m grids, so that 3600 readings were recorded in each grid. These readings were stored in the memory of the instrument and later downloaded to computer for processing and interpretation. Geoplot 3 (Geoscan Research) software was used to process and present the data. Further details are given in Appendix 1.

Reporting

A general site location plan, incorporating the 1:50000 Ordnance Survey (OS) mapping, is shown in Figure 1. Figure 2 is a large scale (1:4000) location plan displaying the processed greyscale magnetometer data. Detailed data plots ('raw' and processed) and an interpretative figure are presented at a scale of 1:1000 in Figures 3 to 8 inclusive.

Further technical information on the equipment used, data processing and survey methodologies is given in Appendix 1 and Appendix 2. Appendix 3 describes the composition and location of the site archive.

The survey methodology, report and any recommendations comply with the Project Design (Harrison 2013) and guidelines outlined by English Heritage (David *et al* 2008) and by the Institute for Archaeologists (IfA 2013). All figures reproduced from Ordnance Survey mapping are with the permission of the controller of Her Majesty's Stationery Office (© Crown copyright).

The figures in this report have been produced following analysis of the data in 'raw' and processed formats and over a range of different display levels. All figures are presented to

most suitably display and interpret the data from this site based on the experience and knowledge of Archaeological Services staff.

4 Results and Discussion

Eller Beck (see Figs 3, 4 and 5)

No anomalies of archaeological potential have been identified by the geophysical survey. Series of east-west orientated parallel linear trend anomalies are caused by former ploughing activity. The vague north-west/south-east aligned band of discrete anomalies (areas of magnetic enhancement), **A**, corresponds closely to a geological fault in the mudstone bedrock at this location (see Fig. 2).

Elsewhere, broad areas of magnetic disturbance within the south and south-east of the site are due to the presence of ferrous material in the perimeter fencing and adjacent golf course. The only other anomalies of note are the occasional ferrous 'spike' anomalies which are caused by small ferrous objects (e.g. nails) within the topsoil. The high magnitude 'spike' within the south of the dataset is due to a larger buried ferrous object.

Waller Hill Beck (see Figs 6, 7 and 8)

The magnetic data collected from the Waller Hill Beck site is characterised by a series of low magnitude, slightly curving, linear trend anomalies, aligned east/west and parallel with the existing field boundaries. These anomalies are due to the medieval and post-medieval agricultural practice of ridge and furrow cultivation. The anomalies are due to the magnetic contrast between the now soil-filled furrows and the former ridges. Towards the south-east of the site two north/south aligned 'speckled' linear trend anomalies are typical of field drains – the 'speckled' effect being caused by the magnetic material (e.g. sand or gravel) within the drain fill. Aside from the ubiquitous ferrous 'spike' anomalies, the only other identified anomalies are low magnitude discrete anomalies which are interpreted as geological in origin, probably resulting from localised variations in the topsoil.

5 Conclusions

No anomalies of archaeological potential have been identified by the geophysical survey. Anomalies indicative of the medieval and post medieval practice of ridge and furrow cultivation have been identified at the Waller Hill Beck site. These anomalies may be of local historical interest but are not thought to be of any archaeological significance. Anomalies due to field drainage and geological variation have also been recorded.

Consequently, on the basis of the geophysical survey, both the Eller Beck site and the Waller Hill Beck site are considered to have a low archaeological potential.

The results and subsequent interpretation of data from geophysical surveys should not be treated as an absolute representation of the underlying archaeological and non-archaeological remains. Confirmation of the presence or absence of archaeological remains can only be achieved by direct investigation of sub-surface deposits.





Fig. 1. Site location




Fig. 2. Survey location showing magnetometer data (1:4000 @ A3)




Fig. 3. Processed greyscale magnetometer data; Eller Beck (1:1000 @ A4) 0 40m




Fig. 4. XY trace plot of minimally processed magnetometer data; Eller Beck (1:1000 @ A4)

0 40m




Fig. 5. Interpretation of magnetometer data; Eller Beck (1:1000 @ A4)

0 40m





Fig. 6. Processed greyscale magnetometer data; Waller Hill Beck (1:1000 @ A3)



Reproduced from the Ordnance Survey mapping with the permission of the Controller of Her Majesty's Stationery Office. © Crown Copyright. Unauthorised reproduction infringes Crown copyright and may lead to prosecution or civil proceedings. Digital data provided by the client, Environment Agency 2013

© ASWYAS 2013.
 Archaeological Services W Y A S
 PO Box 30, Nephshaw Lane South, Morley, LS27 0UG
 Tel: 0113 383 7500 Fax: 0113 383 7501

Fig. 7. XY trace plot of minimally processed magnetometer data; Waller Hill Beck (1:1000 @ A3)

0 40m






Plate 1. General view of Eller Beck survey area, looking north



Plate 2. General view of Waller Hill Beck survey area, looking west



Plate 3. General view of Waller Hill Beck survey area, looking north-east



Plate 4. General view of Waller Hill Beck survey area, looking west

Appendix 1: Magnetic survey - technical information

Magnetic Susceptibility and Soil Magnetism

Iron makes up about 6% of the Earth's crust and is mostly present in soils and rocks as minerals such as maghaemite and haemetite. These minerals have a weak, measurable magnetic property termed magnetic susceptibility. Human activities can redistribute these minerals and change (enhance) others into more magnetic forms so that by measuring the magnetic susceptibility of the topsoil, areas where human occupation or settlement has occurred can be identified by virtue of the attendant increase (enhancement) in magnetic susceptibility. If the enhanced material subsequently comes to fill features, such as ditches or pits, localised isolated and linear magnetic anomalies can result whose presence can be detected by a magnetometer (fluxgate gradiometer).

In general, it is the contrast between the magnetic susceptibility of deposits filling cut features, such as ditches or pits, and the magnetic susceptibility of topsoils, subsoils and rocks into which these features have been cut, which causes the most recognisable responses. This is primarily because there is a tendency for magnetic ferrous compounds to become concentrated in the topsoil, thereby making it more magnetic than the subsoil or the bedrock. Linear features cut into the subsoil or geology, such as ditches, that have been silted up or have been backfilled with topsoil will therefore usually produce a positive magnetic response relative to the background soil levels. Discrete feature, such as pits, can also be detected. The magnetic susceptibility of a soil can also be enhanced by the application of heat and the fermentation and bacterial effects associated with rubbish decomposition. The area of enhancement is usually quite large, mainly due to the tendency of discard areas to extend beyond the limit of the occupation site itself, and spreading by the plough. An advantage of magnetic susceptibility over magnetometry is that a certain amount of occupational activity will cause the same proportional change in susceptibility, however weakly magnetic is the soil, and so does not depend on the magnetic contrast between the topsoil and deeper layers. Susceptibility survey is therefore able to detect areas of occupation even in the absence of cut features. On the other hand susceptibility survey is more vulnerable to the masking effects of layers of colluvium and alluvium as the technique, using the Bartington system, can generally only measure variation in the first 0.15m of ploughsoil.

Types of Magnetic Anomaly

In the majority of instances anomalies are termed 'positive'. This means that they have a positive magnetic value relative to the magnetic background on any given site. However some features can manifest themselves as 'negative' anomalies that, conversely, means that the response is negative relative to the mean magnetic background.

Where it is not possible to give a probable cause of an observed anomaly a '?' is appended.

It should be noted that anomalies interpreted as modern in origin might be caused by features that are present in the topsoil or upper layers of the subsoil. Removal of soil to an archaeological or natural layer can therefore remove the feature causing the anomaly.

The types of response mentioned above can be divided into five main categories that are used in the graphical interpretation of the magnetic data:

Isolated dipolar anomalies (iron spikes)

These responses are typically caused by ferrous material either on the surface or in the topsoil. They cause a rapid variation in the magnetic response giving a characteristic 'spiky' trace. Although ferrous archaeological artefacts could produce this type of response, unless there is supporting evidence for an archaeological interpretation, little emphasis is normally given to such anomalies, as modern ferrous objects are common on rural sites, often being present as a consequence of manuring.

Areas of magnetic disturbance

These responses can have several causes often being associated with burnt material, such as slag waste or brick rubble or other strongly magnetised/fired material. Ferrous structures such as pylons, mesh or barbed wire fencing and buried pipes can also cause the same disturbed response. A modern origin is usually assumed unless there is other supporting information.

Linear trend

This is usually a weak or broad linear anomaly of unknown cause or date. These anomalies are often caused by agricultural activity, either ploughing or land drains being a common cause.

Areas of magnetic enhancement/positive isolated anomalies

Areas of enhanced response are characterised by a general increase in the magnetic background over a localised area whilst discrete anomalies are manifest by an increased response (sometimes only visible on an XY trace plot) on two or three successive traverses. In neither instance is there the intense dipolar response characteristic exhibited by an area of magnetic disturbance or of an 'iron spike' anomaly (see above). These anomalies can be caused by infilled discrete archaeological features such as pits or post-holes or by kilns. They can also be caused by pedological variations or by natural infilled features on certain geologies. Ferrous material in the subsoil can also give a similar response. It can often therefore be very difficult to establish an anthropogenic origin without intrusive investigation or other supporting information.

Linear and curvilinear anomalies

Such anomalies have a variety of origins. They may be caused by agricultural practice (recent ploughing trends, earlier ridge and furrow regimes or land drains), natural geomorphological features such as palaeochannels or by infilled archaeological ditches.

Methodology: Magnetic Susceptibility Survey

There are two methods of measuring the magnetic susceptibility of a soil sample. The first involves the measurement of a given volume of soil, which will include any air and moisture that lies within the sample, and is termed volume specific susceptibility. This method results in a bulk value that is not necessarily fully representative of the constituent components of the sample. For field surveys a Bartington MS2 meter with MS2D field loop is used due to its speed and simplicity. The second technique overcomes this potential problem by taking into account both the volume and mass of a sample and is termed mass specific susceptibility. However, mass specific readings cannot be taken in the field where the bulk properties of a soil are usually unknown and so volume specific readings must be taken. Whilst these values are not fully representative they do allow general comparisons across a site and give a broad indication of susceptibility changes. This is usually enough to assess the susceptibility of a site and evaluate whether enhancement has occurred.

Methodology: Gradiometer Survey

There are two main methods of using the fluxgate gradiometer for commercial evaluations. The first of these is referred to as *magnetic scanning* and requires the operator to visually identify anomalous responses on the instrument display panel whilst covering the site in widely spaced traverses, typically 10m apart. The instrument logger is not used and there is therefore no data collection. Once anomalous responses are identified they are marked in the field with bamboo canes and approximately located on a base plan. This method is usually employed as a means of selecting areas for detailed survey when only a percentage sample of the whole site is to be subject to detailed survey.

The disadvantages of magnetic scanning are that features that produce weak anomalies (less than 2nT) are unlikely to stand out from the magnetic background and so will be difficult to detect. The coarse sampling interval means that discrete features or linear features that are parallel or broadly oblique to the direction of traverse may not be detected. If linear features are suspected in a site then the traverse direction should be perpendicular (or as close as is possible within the physical constraints of the site) to the orientation of the suspected features. The possible drawbacks mentioned above mean that a 'negative' scanning result should be validated by sample detailed magnetic survey (see below).

The second method is referred to as *detailed survey* and employs the use of a sample trigger to automatically take readings at predetermined points, typically at 0.25m intervals, on zig-zag traverses 1m apart. These readings are stored in the memory of the instrument and are later dumped to computer for processing and interpretation. Detailed survey allows the visualisation of weaker anomalies that may not have been detected by magnetic scanning.

During this survey a Bartington Grad601 magnetic gradiometer was used taking readings on the 0.1nT range, at 0.25m intervals on zig-zag traverses 1m apart within 30m by 30m square

grids. The instrument was checked for electronic and mechanical drift at a common point and calibrated as necessary. The drift from zero was not logged.

Data Processing and Presentation

The detailed gradiometer data has been presented in this report in XY trace and greyscale formats. In the former format the data shown is 'raw' with no processing other than grid biasing having been done. The data in the greyscale images has been interpolated and selectively filtered to remove the effects of drift in instrument calibration and other artificial data constructs and to maximise the clarity and interpretability of the archaeological anomalies.

An XY plot presents the data logged on each traverse as a single line with each successive traverse incremented on the Y-axis to produce a 'stacked' plot. A hidden line algorithm has been employed to block out lines behind major 'spikes' and the data has been clipped. The main advantage of this display option is that the full range of data can be viewed, dependent on the clip, so that the 'shape' of individual anomalies can be discerned and potentially archaeological anomalies differentiated from 'iron spikes'. Geoplot 3 software was used to create the XY trace plots.

Geoplot 3 software was used to interpolate the data so that 3600 readings were obtained for each 30m by 30m grid. The same program was used to produce the greyscale images. All greyscale plots are displayed using a linear incremental scale.

Appendix 2: Survey location information

The site grid was laid out using a Trimble VRS differential Global Positioning System (Trimble 5800 model). The accuracy of this equipment is better than 0.01m. The survey grids were then super-imposed onto a base map provided by the client to produce the displayed block locations. However, it should be noted that Ordnance Survey positional accuracy for digital map data has an error of 0.5m for urban and floodplain areas, 1.0m for rural areas and 2.5m for mountain and moorland areas. This potential error must be considered if co-ordinates are measured off hard copies of the mapping rather than using the digital co-ordinates.

Archaeological Services WYAS cannot accept responsibility for errors of fact or opinion resulting from data supplied by a third party.

Appendix 3: Geophysical archive

The geophysical archive comprises:-

- an archive disk containing compressed (WinZip 8) files of the raw data, report text (Microsoft Word 2000), and graphics files (Adobe Illustrator CS2 and AutoCAD 2008) files; and
- a full copy of the report.

At present the archive is held by Archaeological Services WYAS although it is anticipated that it may eventually be lodged with the Archaeology Data Service (ADS). Brief details may also be forwarded for inclusion on the English Heritage Geophysical Survey Database after the contents of the report are deemed to be in the public domain (i.e. available for consultation in the North Yorkshire Historic Environment Record).

Bibliography

ASWYAS, *In Prep. Eller Beck, Waller Hill and Skipton Town Centre, North Yorkshire: Archaeological Watching Brief* Unpublished ASWYAS Report

British Geological Survey, 2013. www.bgs.ac.uk/discoveringGeology/geologyOfBritain/viewer.html . (Viewed December 12th 2013)

David, A., N. Linford, P. Linford and L. Martin. 2008. *Geophysical Survey in Archaeological Field Evaluation: Research and Professional Services Guidelines (2nd edition)* English Heritage

Harrison, D. 2013. *Skipton Flood Mitigation Scheme; Geophysical Survey Project Design* Unpublished ASWYAS Document

Institute for Archaeologists, 2013. *Standard and Guidance for archaeological geophysical survey*. IfA

Soil Survey of England and Wales, 1983. *Soil Survey of England and Wales: Soils of Northern England*, Sheet 1